

Routine Practices and Additional Precautions In All Health Care Settings

Provincial Infectious Diseases
Advisory Committee
(PIDAC)

Ontario Agency for Health Protection and Promotion
Published: August, 2009
Second revision: July 2011



Disclaimer for Best Practice Documents

This document was developed by the Provincial Infectious Diseases Advisory Committee on Infection Prevention and Control (PIDAC-IPC). PIDAC-IPC is a multidisciplinary scientific advisory body who provide evidence-based advice to the Ontario Agency for Health Protection and Promotion (Public Health Ontario) regarding multiple aspects of infectious disease identification, prevention and control. PIDAC-IPC's work is guided by the best available evidence and updated as required. Best Practice documents and tools produced by PIDAC-IPC reflect consensus positions on what the committee deems prudent practice and are made available as a resource to the public health and health care providers.

Suggested Citation:

Ontario Agency for Health Protection and Promotion, Provincial Infectious Diseases Advisory Committee. Routine practices and additional precautions in all health care settings. Toronto, ON: Queen's Printer for Ontario; 2011.

NOTES

This document is intended to provide best practices only. Health care settings are encouraged to work towards these best practices in an effort to improve quality of care.

Provincial Infectious Diseases Advisory Committee (PIDAC)

Ontario Agency for Health Protection and Promotion

www.oahpp.ca

Tel: 647-260-7100

Email: pidac@oahpp.ca

All or part of this report may be reproduced for educational purposes only without permission.

© Queen's Printer for Ontario, 2011

ISBN: 978-1-4435-7027-5

PIDAC-IPC would like to acknowledge the contribution and expertise of the following individuals that participated in the development of this document:

PIDAC-IPC Members:

Dr. Mary Vearncombe, Chair

Medical Director, Infection Prevention and Control,
Microbiology
Sunnybrook Health Sciences Centre, Toronto

Dr. Irene Armstrong

Associate Medical Officer of Health
Toronto Public Health, Toronto

Donna Baker

Manager, Infection Prevention and Control
Bruyère Continuing Care, Elisabeth Bruyere Hospital
Ottawa

Mary Lou Card

Manager, Infection Prevention and Control
London Health Sciences Centre and St. Joseph's Health
Care, London

Judy Dennis

Manager, Infection Prevention and Control
Children's Hospital of Eastern Ontario, Ottawa

Dr. Kevin Katz

Infectious Diseases Specialist and Medical Microbiologist
Medical Director, Infection Prevention and Control
North York General Hospital, Toronto

Dr. Allison McGeer

Director, Infection Control
Mount Sinai Hospital, Toronto

Pat Piaskowski

Network Coordinator
Northwestern Ontario Infection Control Network,
Thunder Bay

Dr. Kathryn Suh

Associate Director, Infection Prevention and Control
Program,
The Ottawa Hospital, Ottawa

Dr. Dick Zoutman

Professor and Chair, Divisions of Medical Microbiology
and of Infectious Diseases
Queen's University, Kingston

Ex-officio members:

Joann Braithwaite

Team Lead, Infectious Diseases and Infection Prevention
and Control
Health Protection and Prevention Branch
Ministry of Health and Long-Term Care, Toronto

Dr. Doug Sider

A/Director, Infectious Disease Prevention and Control
Public Health Ontario, Toronto

Dr. Leon Genesove

Chief Physician, Health Care Unit, Occupational Health
and Safety Branch
Ministry of Labour, Toronto

Public Health Ontario Staff:

Liz Van Horne

Scientific Lead
Public Health Ontario, Toronto

Shirley McDonald

Technical Writer
Public Health Ontario, Toronto

Dr. Maureen Cividino

Occupational Health Physician
Public Health Ontario, Toronto

PIDAC-IPC would also like to acknowledge the contributions to this best practices document in assisting with validation of the French-language translation provided by our Regional Infection Control Network colleagues, Isabelle Langman and Jean-Louis Pitre.

Revisions to *Routine Practices and Additional Precautions in All Health Care Settings*, originally published August 2009:

This document incorporates new information relating to carbapenem resistance in *Enterobacteriaceae*. For more information, see also *Annex A: Screening, Testing and Surveillance for Antibiotic-Resistant Organisms (AROs) in All Health Care Settings*, revised July, 2011.

NOTE: New material is coloured in the text.

Table of Contents

PREAMBLE	6
ABOUT THIS DOCUMENT	6
EVIDENCE FOR RECOMMENDATIONS.....	6
HOW AND WHEN TO USE THIS DOCUMENT	6
ASSUMPTIONS AND BEST PRACTICES FOR INFECTION PREVENTION AND CONTROL	6
I. BACKGROUND	11
MECHANISMS OF TRANSMISSION OF MICROORGANISMS IN HEALTH CARE SETTINGS: THE ‘CHAIN OF TRANSMISSION’ ..	11
PRINCIPLES OF ROUTINE PRACTICES AND RATIONALE	12
PRINCIPLES OF ADDITIONAL PRECAUTIONS AND RATIONALE.....	13
ACCOUNTABILITY OF HEALTH CARE PROVIDERS AND HEALTH CARE ORGANIZATIONS	15
II. BEST PRACTICES	16
ROUTINE PRACTICES.....	16
<i>Elements that Comprise Routine Practices</i>	16
<i>Routine Practices for Visitors</i>	17
<i>Risk Assessment</i>	18
<i>Hand Hygiene</i>	20
<i>Personal Protective Equipment (PPE)</i>	21
<i>Environmental Controls</i>	26
<i>Administrative Controls</i>	30
ADDITIONAL PRECAUTIONS	33
<i>Elements that Comprise Additional Precautions</i>	33
<i>Cohorting</i>	34
<i>Additional Precautions for Visitors</i>	35
<i>Initiation and Discontinuation of Additional Precautions</i>	36
<i>Contact Transmission and Contact Precautions</i>	38
<i>Droplet Transmission and Droplet Precautions</i>	41
<i>Airborne Transmission and Airborne Precautions</i>	43
<i>Combinations of Additional Precautions</i>	47
<i>Protective Environment</i>	48
OCCUPATIONAL HEALTH AND HYGIENE ISSUES.....	49
<i>Post-exposure Follow-up</i>	49
<i>Respiratory Protection Program, Fit-testing and Seal-checking</i>	50
AUDITS OF COMPLIANCE WITH FEEDBACK.....	50
SUMMARY OF RECOMMENDATIONS FOR ROUTINE PRACTICES AND ADDITIONAL PRECAUTIONS IN ALL HEALTH CARE SETTINGS	52
APPENDIX A: RANKING SYSTEM FOR RECOMMENDATIONS.....	61
APPENDIX B: PERFORMING A RISK ASSESSMENT RELATED TO ROUTINE PRACTICES AND ADDITIONAL PRECAUTIONS ..	62
APPENDIX C: DECISION-MAKING RELATED TO ACCOMMODATION AND ADDITIONAL PRECAUTIONS	64
APPENDIX D: TIME REQUIRED FOR AIRBORNE ISOLATION ROOM TO CLEAR <i>M. TUBERCULOSIS</i>	67
APPENDIX E: PIDAC’S ROUTINE PRACTICES FACT SHEET FOR ALL HEALTH CARE SETTINGS.....	68
APPENDIX F: SAMPLE SIGNAGE FOR ENTRANCE TO ROOM OF A PATIENT REQUIRING CONTACT PRECAUTIONS IN ACUTE CARE FACILITIES.....	69
APPENDIX G: SAMPLE SIGNAGE FOR ENTRANCE TO ROOM OF A PATIENT REQUIRING CONTACT PRECAUTIONS IN NON-ACUTE CARE FACILITIES	70
APPENDIX H: SAMPLE SIGNAGE FOR ENTRANCE TO ROOM OF A PATIENT REQUIRING DROPLET PRECAUTIONS IN ALL HEALTH CARE FACILITIES.....	71

APPENDIX I: SAMPLE SIGNAGE FOR ENTRANCE TO ROOM OF A PATIENT REQUIRING DROPLET AND CONTACT PRECAUTIONS IN ACUTE CARE FACILITIES 72

APPENDIX J: SAMPLE SIGNAGE FOR ENTRANCE TO ROOM OF A RESIDENT REQUIRING DROPLET AND CONTACT PRECAUTIONS IN NON-ACUTE CARE FACILITIES..... 73

APPENDIX K: SAMPLE SIGNAGE FOR ENTRANCE TO ROOM OF A PATIENT REQUIRING AIRBORNE PRECAUTIONS IN ALL HEALTH CARE FACILITIES..... 74

APPENDIX L: RECOMMENDED STEPS FOR PUTTING ON AND TAKING OFF PERSONAL PROTECTIVE EQUIPMENT (PPE). 75

APPENDIX M: ADVANTAGES AND DISADVANTAGES OF BARRIER EQUIPMENT 77

APPENDIX N: CLINICAL SYNDROMES AND CONDITIONS WITH REQUIRED LEVEL OF PRECAUTIONS REQUIRED 80

REFERENCES 94

Abbreviations

ABHR	Alcohol-Based Hand Rub
AP	Additional Precautions
ARI	Acute Respiratory Infection
ARO	Antibiotic-Resistant Organism
CCC	Complex Continuing Care
CDI	<i>Clostridium difficile</i> Infection
CRE	Carbapenem-Resistant <i>Enterobacteriaceae</i>
CSA	Canadian Standards Association
DIN	Drug Identification Number (Health Canada)
EMS	Emergency Medical Services
HAI	Health Care-Associated Infection
HEPA	High Efficiency Particulate Air
HSCT	Haematopoietic Stem-cell Transplant
HVAC	Heating, Ventilation and Air Conditioning
ICP	Infection Prevention and Control Professional
LTC	Long-Term Care
MMR	Measles/Mumps/Rubella Vaccine
MOHLTC	Ministry of Health and Long-Term Care (Ontario)
MRSA	Methicillin-Resistant <i>Staphylococcus aureus</i>
NIOSH	National Institute for Occupational Safety and Health (U.S.)
OHA	Ontario Hospital Association
OHSA	<i>Occupational Health and Safety Act</i>
OMA	Ontario Medical Association
PHAC	Public Health Agency of Canada
PIDAC	Provincial Infectious Diseases Advisory Committee
PPE	Personal Protective Equipment
RP	Routine Practices
RP/AP	Routine Practices/Additional Precautions
RSV	Respiratory Syncytial Virus
TB	Tuberculosis
VRE	Vancomycin-Resistant Enterococci

Glossary of Terms

Acute Respiratory Infection (ARI): Any new onset acute respiratory infection that could potentially be spread by the droplet route (either upper or lower respiratory tract), which presents with symptoms of a fever greater than 38°C and a new or worsening cough or shortness of breath (also known as febrile respiratory illness, or FRI). It should be noted that elderly people and people who are immunocompromised may not have a febrile response to a respiratory infection.

Additional Precautions (AP): Additional Precautions (i.e., Contact Precautions, Droplet Precautions and Airborne Precautions) are necessary in addition to Routine Practices for certain pathogens or clinical presentations. These precautions are based on the method of transmission (e.g., contact, droplet, airborne).

Aerosol: Small droplet of moisture that may carry microorganisms. Aerosols may be light enough to remain suspended in the air for short periods of time, allowing inhalation of the microorganism.

Airborne Precautions: Airborne Precautions are used in addition to Routine Practices for clients/patients/residents known or suspected of having an illness transmitted by the airborne route (i.e., by small droplet nuclei that remain suspended in the air and may be inhaled by others).

Alcohol-Based Hand Rub (ABHR): A liquid, gel or foam formulation of alcohol (e.g., ethanol, isopropanol) which is used to reduce the number of microorganisms on hands in clinical situations when the hands are not visibly soiled. ABHRs contain emollients to reduce skin irritation and are less time-consuming to use than washing with soap and water.

Antibiotic-Resistant Organism (ARO): A microorganism that has developed resistance to the action of several antimicrobial agents and that is of special clinical or epidemiological significance.

Barrier Equipment: Personal protective equipment (PPE) used to prevent contamination of skin, mucous membranes or clothing of staff in order to prevent transmission from patient-to-patient. See also, *Personal Protective Equipment*.

Carbapenemase: A class of enzymes that inactivate carbapenem antibiotics by hydrolysing them. In almost all instances, these enzymes hydrolyse not only carbapenem antimicrobials but also first-, second- and third-generation cephalosporins and penicillins (e.g., piperacillin-tazobactam). The genetic information to produce carbapenemases is often located on a mobile genetic element (e.g., plasmid, transposon), which frequently also carries resistance to other classes of antimicrobials, such as fluoroquinolones and aminoglycosides.

Carbapenem-Resistant *Enterobacteriaceae* (CRE): For the purposes of this document, these are *Enterobacteriaceae* that are resistant to carbapenem antimicrobials (e.g., imipenem, meropenem, ertapenem) through the production of carbapenemase.

Chain of Transmission: A model used to understand the infection process.

CHICA-Canada: The Community and Hospital Infection Control Association of Canada, a professional organization of persons engaged in infection prevention and control activities in health care settings. CHICA-Canada members include infection prevention and control professionals from a number of related specialties including nurses, epidemiologists, physicians, microbiology technologists, public health and industry. The CHICA-Canada website is located at: <http://www.chica.org>.

Cleaning: The physical removal of foreign material (e.g., dust, soil) and organic material (e.g., blood, secretions, excretions, microorganisms). Cleaning physically removes rather than kills microorganisms. It is accomplished with water, detergents and mechanical action.

Client/Patient/Resident: Any person receiving care within a health care setting. In this document the term 'patient' refers to client/patient/resident.

Cohorting: The assignment of a geographic area such as a room or a patient care area to two or more clients/patients/residents who are either colonized or infected with the same microorganism, with staffing assignments restricted to the cohorted group of patients. See also, *Staff Cohorting*.

Colonization: The presence and growth of a microorganism in or on a body with growth and multiplication but without tissue invasion or cellular injury or symptoms.

Complex Continuing Care (CCC): Complex continuing care provides continuing, medically complex and specialized services to both young and old, sometimes over extended periods of time. Such care also includes support to families who have palliative or respite care needs.

Contact Precautions: Additional practices to reduce the risk of transmitting infectious agents via contact with an infectious person. Contact Precautions are used in addition to Routine Practices.

Contamination: The presence of an infectious agent on hands or on a surface, such as clothing, gowns, gloves, bedding, toys, surgical instruments, patient care equipment, dressings or other inanimate objects.

Continuum of Care: Across all health care sectors, including settings where emergency (including pre-hospital) care is provided, hospitals, complex continuing care, rehabilitation hospitals, long-term care homes, outpatient clinics, community health centres and clinics, physician offices, dental offices, offices of other health professionals, Public Health and home health care.

Direct Care: Providing hands-on care (e.g., bathing, washing, turning client/patient/resident, changing clothes, continence care, dressing changes, care of open wounds/lesions, toileting).

Disinfectant: A product that is used on medical equipment/devices which results in disinfection of the equipment/device.

Disinfection: The inactivation of disease-producing microorganisms. Disinfection does not destroy bacterial spores. Medical equipment/devices must be cleaned thoroughly before effective disinfection can take place.

Droplet Precautions: Droplet Precautions are used in addition to Routine Practices for clients/patients/residents known or suspected of having an infection that can be transmitted by large infectious droplets.

Environment of the Client/Patient/Resident: The immediate space around a client/patient/resident that may be touched by the client/patient/resident and may also be touched by the health care provider when providing care. The client/patient/resident environment includes equipment, medical devices, furniture (e.g., bed, chair, bedside table), telephone, privacy curtains, personal belongings (e.g., clothes, books) and the bathroom that the client/patient/resident uses. In a multi-bed room, the client/patient/resident environment is the area inside the individual's curtain. In an ambulatory setting, the client/patient/resident environment is the area that may come into contact with the client/patient/resident within their cubicle. In a nursery/neonatal setting, the patient environment is the isolette or bassinet and equipment outside the isolette/bassinet that is used for the infant. See also, *Health Care Environment*.

Extended Spectrum Beta-Lactamase (ESBL): Extended-spectrum β -lactamases are enzymes that may be produced by some strains of *Enterobacteriaceae* that hydrolyse all cephalosporins, including third-generation cephalosporins such as cefotaxime, ceftriaxone and ceftazidime, as well as aztreonam.

Eye Protection: A device that covers the eyes and is used by health care providers to protect the eyes when it is anticipated that a procedure or care activity is likely to generate splashes or sprays of blood, body fluids, secretions or excretions, or within two metres of a coughing client/patient/resident. Eye protection includes safety glasses, safety goggles, face shields and visors.

Facial Protection: Personal protective equipment that protect the mucous membranes of the eyes, nose and mouth from splashes or sprays of blood, body fluids, secretions or excretions. Facial protection may include a mask or respirator in conjunction with eye protection, or a face shield that covers eyes, nose and mouth.

Fit-Check: See *Seal-Check*

Fit-Test: A qualitative or quantitative method to evaluate the fit of a specific make, model and size of respirator on an individual. Fit-testing is to be done periodically, at least every two years and whenever there is a change in respirator face piece or the user's physical condition which could affect the respirator fit.⁴

Hand Care Program: A hand care program for staff is a key component of hand hygiene and includes hand care assessment, staff education, Occupational Health assessment if skin integrity is an issue, provision of hand moisturizing products and provision of alcohol-based hand rub that contains an emollient. For more information about implementing a hand care program, refer to *Best Practices for Hand Hygiene in All Health Care Settings*⁵, available at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/hand-hygiene.html>.

Hand Hygiene: A general term referring to any action of hand cleaning. Hand hygiene relates to the removal of visible soil and removal or killing of transient microorganisms from the hands. Hand hygiene may be accomplished using soap and running water or an alcohol-based hand rub. Hand hygiene also includes surgical hand antisepsis.

Hand Washing: The physical removal of microorganisms from the hands using soap (plain or antimicrobial) and running water.

Health Care-Associated Infection (HAI): A term relating to an infection that is acquired during the delivery of health care (also known as *nosocomial infection*).

Health Care Environment: People and items which make up the care environment (e.g., objects, medical equipment, staff, clients/patients/residents) of a hospital, clinic or ambulatory setting, outside the immediate environment of the client/patient/resident. See also, *Environment of the Client/Patient/Resident*.

Health Care Facility: A set of physical infrastructure elements supporting the delivery of health-related services. A health care facility does not include a client/patient/resident's home or physician/dentist/other health offices where health care may be provided.

Health Care Provider: Any person delivering care to a client/patient/resident. This includes, but is not limited to, the following: emergency service workers, physicians, dentists, nurses, respiratory therapists and other health professionals, personal support workers, clinical instructors, students and home health care workers. In some non-acute settings, volunteers might provide care and would be included as a health care provider. See also, *Staff*.

Health Care Setting: Any location where health care is provided, including settings where emergency care is provided, hospitals, complex continuing care, rehabilitation hospitals, long-term care homes, mental health facilities, outpatient clinics, community health centres and clinics, physician offices, dental offices, offices of other health professionals and home health care.

HEPA Filter: High efficiency particulate air filter with an efficiency of 99.97% in the removal of airborne particles 0.3 microns or larger in diameter.⁶

Hospital-Grade Disinfectant: A disinfectant that has a drug identification number (DIN) from Health Canada indicating its approval for use in Canadian hospitals.

Infection: The entry and multiplication of an infectious agent in the tissues of the host. Asymptomatic or sub-clinical infection is an infectious process running a course similar to that of clinical disease but below the threshold of clinical symptoms. Symptomatic or clinical infection is one resulting in clinical signs and symptoms (disease).

Infection Prevention and Control: Evidence-based practices and procedures that, when applied consistently in health care settings, can prevent or reduce the risk of transmission of microorganisms to health care providers, other clients/patients/residents and visitors.

Infection Prevention and Control Professional(s) (ICPs): Trained individual(s) responsible for a health care setting's infection prevention and control activities. In Ontario an ICP must receive a minimum of 80 hours of instruction in a CHICA-Canada endorsed infection control program within six months of entering the role and must acquire and maintain Certification in Infection Control (CIC) when eligible.

Infectious Agent: A microorganism, i.e., a bacterium, fungus, parasite, virus or prion, which is capable of invading body tissues and multiplying.

Long-Term Care (LTC): A broad range of personal care, support and health services provided to people who have limitations that prevent them from full participation in the activities of daily living. The people who use long-term care services are usually the elderly, people with disabilities and people who have a chronic or prolonged illness.

Mask: A device that covers the nose and mouth, is secured in the back and is used by health care providers to protect the mucous membranes of the nose and mouth.

Methicillin-Resistant *Staphylococcus aureus* (MRSA): MRSA is a strain of *Staphylococcus aureus* that has a minimal inhibitory concentration (MIC) to oxacillin of ≥ 4 mcg/ml and contains the *mecA* gene coding for penicillin-binding protein 2a (PBP 2a). MRSA is resistant to all of the beta-lactam classes of antibiotics, such as penicillins, penicillinase-resistant penicillins (e.g., cloxacillin) and cephalosporins.

N95 Respirator: A personal protective device that is worn on the face and covers the nose and mouth to reduce the wearer's risk of inhaling airborne particles. A NIOSH-certified N95 respirator filters particles one micron in size, has 95% filter efficiency and provides a tight facial seal with less than 10% leak.^{7,8}

Occupational Health: Health services in the workplace provided by trained occupational health nurses and physicians.

Personal Protective Equipment (PPE): Clothing or equipment worn by staff for personal protection against hazards. See also, *Barrier Equipment*.

Point-of-Care: The place where three elements occur together: the client/patient/resident, the health care provider and care or treatment involving client/patient/resident contact.

Pre-Hospital Care: Acute emergency client/patient/resident assessment and care delivered in an uncontrolled environment by designated practitioners, performing delegated medical acts at the entry to the health care continuum.

Provincial Infectious Diseases Advisory Committee (PIDAC): A multidisciplinary scientific advisory body who provide evidence-based advice to the Ontario Agency for Health Protection and Promotion (Public Health Ontario) regarding multiple aspects of infectious disease identification, prevention and control. More information is available at: <http://www.oahpp.ca/services/pidac/index.html>.

Public Health Agency of Canada (PHAC): A national agency which promotes improvement in the health status of Canadians through public health action and the development of national guidelines. The PHAC website is located at: http://www.phac-aspc.gc.ca/new_e.html.

Regional Infection Control Networks (RICN): The RICN of Ontario coordinate and integrate resources related to the prevention, surveillance and control of infectious diseases across all health care sectors and for all health care providers, promoting a common approach to infection prevention and control and utilization of best-practices within the region. There are 14 regional networks in Ontario. More information is available at: <http://www.ricn.on.ca>.

Respirator: See *N95 respirator*.

Respiratory Etiquette: Personal practices that help prevent the spread of bacteria and viruses that cause acute respiratory infections (e.g., covering the mouth when coughing, care when disposing of tissues).

Risk Assessment: An evaluation of the interaction of the health care provider, the client/patient/resident and the client/patient/resident environment to assess and analyze the potential for exposure to infectious disease.

Routine Practices (RP): The system of infection prevention and control practices recommended by the Public Health Agency of Canada to be used with all clients/patients/residents during all care to prevent and control transmission of microorganisms in health care settings.

Seal-Check: A procedure that the health care provider must perform each time an N95 respirator is worn to ensure it fits the wearer's face correctly to provide adequate respiratory protection. The health care provider is to receive training on how to perform a seal-check correctly.⁴

Sharps: Objects capable of causing punctures or cuts (e.g., needles, syringes, blades, clinical glass).

Staff: Anyone conducting activities in settings where health care is provided, including but not limited to, health care providers. See also, *Health Care Providers*.

Staff Cohorting: The practice of assigning specified health care providers to care only for clients/patients/residents known to be colonized or infected with the same microorganism. These health care providers would not participate in the care of clients/patients/residents who are not colonized or infected with that microorganism. See also, *Cohorting*.

Terminal Cleaning: The cleaning of a client/patient/resident room or bed space following discharge or transfer of the client/patient/resident, in order to remove contaminating microorganisms that might be acquired by subsequent occupants. In some instances, terminal cleaning might be used once some types of Additional Precautions have been discontinued. Terminal cleaning methods vary, but usually include removing all detachable objects in the room, cleaning lighting and air duct surfaces in the ceiling, and cleaning everything downward to the floor. Items removed from the room are disinfected before being returned to the room. Refer to *Best Practices for Environmental Cleaning in All Health Care Settings*⁹ for more information about terminal cleaning.

Vancomycin-Resistant Enterococci (VRE): VRE are strains of *Enterococcus faecium* or *Enterococcus faecalis* that have a minimal inhibitory concentration (MIC) to vancomycin of ≥ 32 mcg/ml. and/or contain the resistance genes vanA or vanB.

Preamble

About This Document

This document outlines the practice of Routine Practices and Additional Precautions (RP/AP) in health care settings across the continuum of care (see below) including, but not limited to, pre-hospital care, acute care, complex continuing care, rehabilitation facilities, long-term care, chronic care, ambulatory care and home health care.

The goal of Routine Practices and Additional Precautions is to reduce the risk of transmission of microorganisms in health care settings through:

- understanding the concepts of the chain of transmission;
- understanding the concepts and application of Routine Practices (RP);
- understanding barriers and enablers that affect compliance with Routine Practices;
- knowing why and when to use Additional Precautions (AP); and
- using, applying and removing personal protective equipment correctly when indicated for the protection of the client/patient/resident or the staff member.

For recommendations in this document:

- **'shall'** indicates mandatory requirements based on legislated requirements or national standards (i.e., Canadian Standards Association - CSA)
- **'must'** indicates best practice, i.e., the minimum standard based on current recommendations in the medical literature;
- **'should'** indicates a recommendation or that which is advised but not mandatory; and
- **'may'** indicates an advisory or optional statement.

Evidence for Recommendations

The best practices in this document reflect the best evidence and expert opinion available at the time of writing. As new information becomes available, this document will be reviewed and updated. [Refer to Appendix A](#), 'Ranking System for Recommendations', for grading system used for recommendations.

How and When to Use This Document

The Routine Practices and Additional Precautions set out in this document must be practiced in all settings where health care is provided, across the continuum of health care. This includes settings where emergency (including pre-hospital) care is provided, hospitals, complex continuing care facilities, rehabilitation facilities, long-term care homes, outpatient clinics, community health centres and clinics, physician offices, dental offices, offices of other health professionals, public health and home health care.

Assumptions and Best Practices for Infection Prevention and Control

The best practices in this document are based on the assumption that health care settings in Ontario already have basic infection prevention and control systems in place.¹⁰ These settings should work with organizations that have infection prevention and control expertise, such as academic health science centres, regional infection control networks, public health units that have professional staff certified in infection prevention and control and local infection prevention and control associations (e.g., Community and Hospital Infection Control Association (CHICA) – Canada chapters), to develop evidence-based programs.

In addition to the above general assumption about basic infection prevention and control, these best practices are based on the following additional assumptions and principles:

1. Adequate resources are devoted to infection prevention and control in all health care settings. See *Best Practices for Infection Prevention and Control Programs in Ontario*,¹⁰ available from the Public Health Ontario (PHO) website at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/infection-prevention-and-control-programs-in-ontario.html>
2. Programs are in place in all health care settings that promote good hand hygiene practices and ensure adherence to standards for hand hygiene. See *Best Practices for Hand Hygiene in All Health Care Settings*,¹¹ available from the PHO website at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/hand-hygiene.html>.

See also Ontario's hand hygiene improvement program, *Just Clean Your Hands*, available at: <http://www.health.gov.on.ca/en/ms/handhygiene/>

3. Adequate resources are devoted to Environmental Services/Housekeeping in all health care settings that include written procedures for cleaning and disinfection of client/patient/resident rooms and equipment; education of new cleaning staff and continuing education of all cleaning staff; and ongoing review of procedures. See *Best Practices for Environmental Cleaning in All Health Care Settings*,⁹ available from the PHO website at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/environmental-cleaning-for-prevention-and-control-of-infections.html>
4. Programs are in place in all health care settings that ensure effective disinfection and sterilization of used medical equipment according to *Best Practices for Cleaning, Disinfection and Sterilization in All Health Care Settings*,¹² available from the PHO website at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/cleaning-disinfection-and-sterilization.html>
5. Regular education (including orientation and continuing education) and support to help staff consistently implement appropriate infection prevention and control practices is provided in all health care settings.

Effective education programs emphasize:

- the risks associated with infectious diseases, including acute respiratory illness and gastroenteritis
- hand hygiene, including the use of alcohol-based hand rubs and hand washing
- assessment of the risk of infection transmission and the appropriate use of personal protective equipment (PPE), including safe application, removal and disposal as defined in this document
- principles and components of Routine Practices as well as additional transmission-based precautions, as set out in this document
- appropriate cleaning and/or disinfection of health care equipment, supplies and surfaces or items in the health care environment
- individual staff responsibility for keeping clients/patients/residents, themselves and co-workers safe
- collaboration between professionals involved in occupational health and infection prevention and control.

NOTE: Education programs should be flexible enough to meet the diverse needs of the range of health care providers and other staff who work in the health care setting. The local public health unit and regional infection control networks may be a resource and can provide assistance in developing and providing education programs for community settings.

6. Collaboration between professionals involved in occupational health and infection prevention and control is promoted in all health care settings to implement and maintain appropriate infection prevention and control standards that protect workers.
7. There are effective working relationships between the health care setting and the local public health unit. Clear lines of communication are maintained and public health is contacted for information and advice as required and the obligations (under the *Health Protection and Promotion Act*, R.S.O. 1990, c.H.7)¹³ to report reportable and communicable diseases is fulfilled. Public health provides regular aggregate reports of outbreaks of any infectious diseases in facilities and/or in the community to all health care settings.
8. Access to ongoing infection prevention and control advice and guidance to support staff and resolve differences is available to the health care setting.
9. There are established procedures for receiving and responding appropriately to all international, national, regional and local health advisories in all health care settings. Health advisories are communicated promptly to all staff responsible for case finding/surveillance and regular updates are provided. Current advisories are available from local public health units, the MOHLTC, Health Canada and Public Health Agency of Canada websites and local regional infection prevention and control networks.
10. Where applicable, there is a process for evaluating personal protective equipment (PPE) in the health care setting, to ensure it meets quality standards.
11. There is regular assessment of the effectiveness of the infection prevention and control program and its impact on practices in the health care setting. The information is used to further refine the program.¹⁰
12. The Ministry of Health and Long-Term Care's Long-Term Care Home Compliance and Enforcement Program requirements shall be met. Specific legislative requirements for long-term care providers may be found in The *Long-Term Care Homes Act, 2007* available at:

http://www.e-laws.gov.on.ca/html/statutes/english/elaws_statutes_07108_e.htm

This act contains specific requirements respecting infection control including:

- S.76(2) that every licensee must ensure that staff have received training in infection prevention and control;
- S.86 requiring every licensee to ensure there is a program of infection prevention and control

More detailed requirements of the infection prevention and control program are found in Regulation 79/10 available at: http://www.e-laws.gov.on.ca/html/regs/english/elaws_regs_100079_e.htm#BK285

In addition, all long-term care providers shall comply with all requirements outlined in the MOHLTC's '*Long-Term Care Homes Program Manual*',¹⁴ which is the core text governing the operation of long-term care homes in the province of Ontario. This manual contains policies, standards and norms covering various aspects of the LTC Homes Program such as:

- Risk Management, including:
 - infection control
 - health and safety
 - internal and external disaster planning
 - monitoring, evaluating and improving quality
- Environmental Services, including:
 - waste management
 - pest control
 - housekeeping services
 - laundry services

- maintenance services
- Education, including:
 - orientation
 - ongoing inservice education
 - mandatory education programs
- The Program Manual may be accessed at:

http://www.health.gov.on.ca/english/providers/pub/manuals/ltc_homes/ltc_homes_mn.html#full.
- For more information, please contact your local Ministry of Health Service Area Office. A list of these offices may be found at:

<http://www.infogo.gov.on.ca/infogo/office.do?actionType=telephonedirectory&infoType=telephone&unitId=UNT0028407&locale=en>.

13. Occupational Health and Safety requirements shall be met:

- Health care facilities are required to comply with applicable provisions of the *Occupational Health and Safety Act* (OHSA), R.S.O. 1990, c.0.1 and its Regulations.¹⁵ Employers, supervisors and workers have rights, duties and obligations under the OHSA. To see what the specific requirements are under the OHSA go to: http://www.e-laws.gov.on.ca/html/statutes/english/elaws_statutes_90o01_e.htm
- The *Occupational Health and Safety Act* places duties on many different categories of individuals associated with workplaces, such as employers, constructors, supervisors, owners, suppliers, licensees, officers of a corporation and workers. A guide to the requirements of the *Occupational Health and Safety Act* may be found at: <http://www.labour.gov.on.ca/english/hs/ohsaguide/index.html>
- Specific requirements for certain health care and residential facilities may be found in the *Regulation for Health Care and Residential Facilities*. Go to: http://www.e-laws.gov.on.ca/html/regs/english/elaws_regs_930067_e.htm
- In addition, the OHSA section 25(2)(h), the ‘general duty clause’, requires an employer to take every precaution reasonable in the circumstances for the protection of a worker. There is a general duty for an employer to establish written measures and procedures for the health and safety of workers, in consultation with the joint health and safety committee or health and safety representative, if any. Such measures and procedures may include, but are not limited to, the following:
 - safe work practices
 - safe working conditions
 - proper hygiene practices and the use of hygiene facilities
 - the control of infections.
- At least once a year the measures and procedures for the health and safety of workers shall be reviewed and revised in the light of current knowledge and practice. The employer, in consultation with the joint health and safety committee or health and safety representative, if any, shall develop, establish and provide training and educational programs in health and safety measures and procedures for workers that are relevant to the workers’ work.
- A worker who is required by his or her employer or by the *Regulation for Health Care and Residential Facilities* to wear or use any protective clothing, equipment or device shall be instructed and trained in its care, use and limitations before wearing or using it for the first time and at regular intervals thereafter and the worker shall participate in such instruction and training. The employer is reminded of the need to be able to demonstrate training, and is therefore encouraged to document the workers trained, the dates training was conducted, and materials covered during training. Under the *Occupational Health and Safety Act*, a worker must work in compliance with the Act and its

regulations, and use or wear any equipment, protective devices or clothing required by the employer.

- For more information, please contact your local Ministry of Labour office. A list of local Ministry of Labour offices in Ontario may be found at <http://www.labour.gov.on.ca/>

Routine Practices and Additional Precautions in All Health Care Settings

I. Background


TERMS USED IN THIS DOCUMENT (see glossary for details and examples)

Health Care Provider: Any person delivering care to a client/patient/resident

Staff: Anyone conducting activities within a health care setting (includes health care providers)


Mechanisms of Transmission of Microorganisms in Health Care Settings: The 'Chain of Transmission'

The transmission of microorganisms and subsequent infection within a health care setting may be likened to a 'chain', with each link in the chain representing a factor related to the spread of microorganisms. Transmission does not take place unless all six of the elements in the chain of transmission are present (see [Figure 1](#)). By eliminating any of the six links, or 'breaking the chain', transmission does not occur (see [Figure 2](#)).



Transmission occurs when the agent, in the reservoir, exits the reservoir through a portal of exit, travels via a mode of transmission and gains entry through a portal of entry to a susceptible host.

Figure 1: The Chain of Transmission



Transmission may be interrupted when:

- **the agent is eliminated or inactivated or cannot exit the reservoir;**
- **portals of exit are eliminated through safe practices;**
- **transmission between objects or people does not occur due to barriers and/or safe practices;**
- **portals of entry are protected; and/or**
- **hosts are not susceptible**

Figure 2: Breaking the Chain of Transmission

Principles of Routine Practices and Rationale

Routine Practices are based on the premise that all clients/patients/residents are *potentially* infectious, even when asymptomatic, and that the same safe standards of practice should be used **routinely** with **all** clients/patients/residents to prevent exposure to blood, body fluids, secretions, excretions, mucous membranes, non-intact skin or soiled items and to prevent the spread of microorganisms.

The consistent and appropriate use of Routine Practices by all health care providers with all patient encounters will lessen microbial transmission in the health care setting and reduce the need for Additional Precautions.

The risk of transmission of microorganisms involves factors related to the microbe, the source client/patient/resident, the health care environment and the new host.⁷

Health care providers must assess the risk of exposure to blood, body fluids and non-intact skin and identify the strategies that will decrease exposure risk and prevent the transmission of microorganisms. This risk assessment followed by the implementation of Routine Practices to reduce or remove risk should be incorporated into the culture of each health care setting and into the daily practice of each health care provider. The goals of Routine Practices are listed in [Figure 3](#).

Health care providers must assess the risk of exposure to blood, body fluids and non-intact skin and identify the strategies that will decrease exposure risk and prevent the transmission of microorganisms.




Figure 3: Goals of Routine Practices

Principles of Additional Precautions and Rationale

Additional Precautions are used in addition to Routine Practices for clients/patients/residents known or suspected to be infected or colonized with certain microorganisms to interrupt transmission. Refer to [Appendix N](#) for a list of microorganisms/diseases that require Additional Precautions.

Additional Precautions include the use of barriers, such as PPE, and additional environmental controls that are put in place for encounters with the client/patient/resident or their immediate environment. In some instances, specialized engineering controls may be required (e.g., negative pressure room for a patient with Tuberculosis) or enhanced cleaning protocols for the client/patient/resident environment (e.g., *Clostridium difficile* – *C. difficile*, vancomycin-resistant enterococci - VRE).

The application of Additional Precautions may differ depending on the health care setting and the needs of the client/patient/resident, particularly in long-term care and the community. More information about Additional Precautions is available in Section II.

Staff in all health care settings must follow Routine Practices and Additional Precautions and facilities must implement a program that includes:

- a) written policies and procedures that include risk assessment;
- b) staff education and training in indications and techniques for Routine Practices and Additional Precautions, including hand hygiene;

- c) a program to measure compliance with Routine Practices and Additional Precautions, including hand hygiene;
- d) sufficient and easily accessible personal protective equipment (e.g., gloves, masks, eye protection, gowns) available for health care providers and other staff who are exposed to blood and body substances with education and training in their use;
- e) healthy workplace policies including a sharps injury prevention program;16 staff immunization program; requirement for staff to remain home if ill with an infection which may be transmitted to clients/patients/residents or other staff; and promotion of respiratory etiquette for clients/patients/residents and staff; and
- f) appropriate environmental controls that reduce the risks of transmission of microorganisms.

Successful implementation of Routine Practices and Additional Precautions (RP/AP) requires the support of senior administration. See [Figure 4](#) for components required for the successful implementation of Routine Practices and Additional Precautions in health care facilities.




Figure 4: Components Required When Implementing Routine Practices and Additional Precautions

Accountability of Health Care Providers and Health Care Organizations

Adherence to recommended infection prevention and control practices decreases transmission of microorganisms in health care settings.¹⁷⁻²¹ Despite this, there are numerous studies on the behaviour of health care providers that show poor compliance with hand hygiene²²⁻²⁴ and the use of protective barrier equipment,²⁵⁻²⁷ placing both staff and clients/patients/residents at risk.

Organizations have a responsibility to have systems in place with established procedures that enable compliance with Hand Hygiene, Routine Practices and Additional Precautions. Both the employer and the employee have duties under the *Occupational Health and Safety Act*¹⁵:

- a) 'An employer shall ensure that the equipment, materials and protective devices as prescribed are provided' [S. 25(1)(a)] and 'the equipment, materials and protective devices provided by the employer are maintained in good condition' [S. 25(1)(b)];
- b) 'A worker shall use or wear the equipment, protective devices or clothing that his employer requires to be used or worn' [S. 28(1)(b)] and 'a worker shall report to his or her employer or supervisor the absence of or defect in any equipment or protective device of which the worker is aware and which may endanger himself, herself or another worker' [S. 28(1)(c)].

Preventing transmission of microorganisms to other clients/patients/residents is a patient safety issue, and preventing transmission to staff is an occupational health and safety issue. Health care providers are accountable to practice safely in a manner that protects clients/patients/residents and themselves by following established organizational infection prevention and control policies and procedures.

The consistent and appropriate use of Routine Practices by all health care providers will lessen microbial transmission in the health care setting and reduce the need for Additional Precautions.

II. Best Practices

Routine Practices

Routine Practices refer to infection prevention and control practices to be used with all clients/patients/residents during all care, to prevent and control transmission of microorganisms in all health care settings. Routine Practices must be incorporated into the culture of each health care setting and into the daily practice of each health care provider.


Routine Practices must be incorporated into the culture of each health care setting and into the daily practice of each health care provider.

ELEMENTS THAT COMPRISE ROUTINE PRACTICES

The basic elements of Routine Practices are listed in Box 1 and include:

- a) **risk assessment** of the client/patient/resident and the health care provider's interaction with the client/patient/resident;
- b) **hand hygiene** to be performed with an alcohol-based hand rub or with soap and water before and after contact with a client/patient/resident or their environment, before invasive/aseptic procedures and after contact with body fluids;
 - Refer to *Best Practices for Hand Hygiene in All Health Care Settings*¹¹, available on the PHO website at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/hand-hygiene.html>
 - Refer to the *Just Clean Your Hands* program for more information about hand hygiene; available at: <http://www.health.gov.on.ca/en/ms/handhygiene/>
- c) environmental controls, including:
 - i) appropriate **placement and bed spacing**, such as single room and private toileting facilities for clients/patients/residents who soil the environment;
 - ii) **cleaning of equipment** that is being used by more than one client/patient/resident between uses according to the recommendations found in *Best Practices For Cleaning, Disinfection and Sterilization in All Health Care Settings*,¹² available at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/cleaning-disinfection-and-sterilization.html>
 - iii) **cleaning of the health care environment**, including safe handling of soiled linen and waste (e.g., sharps) to prevent exposure and transmission to others, as detailed in *Best Practices for Environmental Cleaning in All Health Care Settings*, available on the PHO website at:⁹ <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/environmental-cleaning-for-prevention-and-control-of-infections.html>; and
 - iv) **engineering controls**, such as well-maintained heating, ventilation and air conditioning (HVAC) systems with sufficient air changes per hour; and
 - v) **point-of-care sharps containers**, hand hygiene product dispensers and adequate dedicated hand wash sinks;

- d) administrative controls including:
 - i) **policies and procedures** to ensure that staff are able to deal effectively with transmission risks associated with infectious illnesses;
 - ii) **staff education** to heighten awareness of infectious diseases, their mode of transmission and prevention of transmission;
 - iii) **healthy workplace policies** that exclude staff from working when ill with a communicable disease that would put clients/patients/residents and colleagues at risk;
 - iv) **immunization programs** for staff and for clients/patients/residents where applicable;
 - v) **respiratory etiquette** for both staff and clients/patients/residents;
 - vi) **monitoring of compliance with feedback** is built into the program to measure compliance with Routine Practices, including hand hygiene; and
 - vii) **sufficient staffing levels** to enable health care providers to comply with infection prevention and control policies and procedures;
- e) **sufficient, easily accessible and appropriate barrier equipment** (i.e., personal protective equipment) to prevent health care provider contact with blood, body fluids, secretions, excretions, non-intact skin or mucous membranes.



ROUTINE PRACTICES FOR VISITORS

Although visitors are less likely to transmit infection in the health care setting than staff, they should receive instruction regarding specific facility control measures before they visit a client/patient/resident, to ensure compliance with established practices:

- a) visitors should not enter the health care setting if they are sick or unable to comply with hand hygiene and other precautions that might be required;
- b) hand hygiene before and after visiting should be emphasized; and
- c) if barrier equipment is required by the visitor, this should be accompanied by instruction in its correct application, use and disposal.

Instructional materials may be provided to visitors on recommended hand hygiene and respiratory etiquette practices.

RISK ASSESSMENT

The first step in the effective use of Routine Practices is to perform a risk assessment. A risk assessment must be done **before each interaction** with a client/patient/resident or their environment in order to determine which interventions are required to prevent transmission during the interaction,²⁸ because the client/patient/resident's status can change.

The risk assessment process will be a dynamic one, based on continuing changes in information as care progresses, thus must be done before each interaction with a client/patient/resident.

Assessing Risk of Transmission

The risk of transmission of microorganisms between individuals involves factors related to:

- a) the client/patient/resident infection status (including colonization);
- b) the characteristics of the client/patient/resident;
- c) the type of care activities to be performed;
- d) the resources available for control; and
- e) the health care provider immune status.⁷







Table 1 lists factors affecting the risk of transmission of microorganisms in health care settings. The health care provider must perform a risk assessment of each task or interaction that includes:

- a) assessing the risk of:
 - i) contamination of skin or clothing by microorganisms in the client/patient/resident environment;
 - ii) exposure to blood, body fluids, secretions, excretions, tissues;
 - iii) exposure to non-intact skin;
 - iv) exposure to mucous membranes; and
 - v) exposure to contaminated equipment or surfaces; and
- b) recognition of symptoms of infection (e.g., syndromic surveillance).¹⁸ See Box 9 for a list of clinical syndromes requiring the use of PPE and other controls pending diagnosis.

Where there is a risk of transmission of infection based on the risk assessment, appropriate controls must be put into place and appropriate PPE must be used to protect the health care provider, other staff and clients/patients/residents until a definitive diagnosis may be made. For example:

- If a client/patient/resident has uncontained diarrhea, barrier equipment such as gloves and a gown should be considered when changing the bed sheets, to prevent contamination of hands and clothing;
 - If the client/patient/resident is soiling the environment outside of the immediate bed area, a single room is preferable to limit transmission to other clients/patients/residents;
 - Use avoidance procedures that **minimize contact with droplets** (e.g., sitting next to, rather than in front of, a coughing client/patient/resident when taking a history or conducting an examination).
- Refer to Appendix B, 'Performing a Risk Assessment Related to Routine Practices and Additional Precautions', for more information related to risk assessment.

TABLE 1: FACTORS AFFECTING RISK OF TRANSMISSION OF MICROORGANISMS IN A HEALTH CARE SETTING

Higher risk of transmission is associated with:	Link in chain of transmission
<p>Microorganism/Infectious Agent</p> <ul style="list-style-type: none"> • Presence of a large amount of the infectious agent • Low infective dose required for infection (i.e., high infectivity) • High pathogenicity/virulence • Airborne-spread • Able to survive in the environment • Able to colonize invasive devices • Able to exist in an asymptomatic/carrier state 	
<p>Source Client/Patient/Resident</p> <ul style="list-style-type: none"> • Incontinent of stool and stool not contained by incontinence products • Draining skin lesions or wounds not contained by dressings • Copious uncontrolled respiratory secretions • Inability to comply with hygienic practices and infection prevention and control precautions • Patient in intensive care unit or requiring extensive hands-on care 	 
<p>Environment</p> <ul style="list-style-type: none"> • Inadequate cleaning • Shared care equipment without cleaning between clients/patients/residents • Crowded facilities • Shared facilities, such as multi-bed rooms (e.g., toilets, sinks, baths) • High patient-nurse ratio 	
<p>Susceptible Host</p> <ul style="list-style-type: none"> • Patient in intensive care unit or requiring extensive hands-on care • Patient has invasive procedures or devices • Non-intact skin (client/patient/resident or staff) • Debilitated, severe underlying disease • Extremes of age • Recent antibiotic therapy • Immunosuppression • Lack of appropriate immunization • Inadequately educated, trained or non-compliant staff 	 

Adapted from: ‘Routine Practices and Additional Precautions for Preventing the Transmission of Infection in Health Care’; Health Canada, CCDR July 1999; Vol. 25 Supplement 4: p.19

HAND HYGIENE

Hand hygiene is considered the most important and effective infection prevention and control measure to prevent the spread of health care-associated infections. In order to implement a comprehensive hand hygiene program in a health care facility, refer to:

- The *Just Clean Your Hands* hand hygiene improvement program for hospitals and long-term care,²⁹ available at: <http://www.health.gov.on.ca/en/ms/handhygiene/>
- *Best Practices for Hand Hygiene in All Health Care Settings*,¹¹ available on the PHO website at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/hand-hygiene.html>
- The World Health Organization's *Clean Care is Safer Care* hand hygiene campaign is available at: <http://www.who.int/gpsc/en/>

Hand Hygiene Program

In spite of its importance in the prevention of health care-associated infections (HAIs), compliance with hand hygiene by health care providers has been, and continues to be, unacceptably low at 20% to 50%.^{19, 30, 31} It has been shown that a multifaceted, multidisciplinary, facility-wide hand hygiene program, which includes demonstrable administrative leadership, education, champions and environmental enablers can be effective at reducing the incidence of HAIs.¹⁹

All health care settings must implement a comprehensive hand hygiene program that incorporates the following elements¹⁰:

- a) the program is multifaceted and multidisciplinary to provide leadership and decision-making;
- b) hand hygiene agents are available at point-of-care in all health care settings; hand hygiene agents must be dispensed in disposable containers and must not be 'topped up';
- c) education is given to health care providers about when and how to clean their hands; and
- d) a hand care program to maintain skin integrity in collaboration with Occupational Health.

Health care facilities must also include¹⁰:

- a) senior and middle management support and commitment to make hand hygiene an organizational priority and address non-compliance;
- b) environmental changes and system supports, including alcohol-based hand rub at the point-of-care and a hand care program;
- c) ongoing auditing and observation of hand hygiene practices, with feedback to health care providers;
- d) client/patient/resident engagement; and
- e) opinion leaders and champions modeling the right behaviour.

Alcohol-based Hand Rub (ABHR)

To make it possible for health care providers to clean their hands at the right time, alcohol-based hand rub (ABHR) or a hand hygiene sink must be provided at the point-of-care, where busy health care providers can clean their hands without leaving the client/patient/resident.³² ABHRs are the preferred method to routinely decontaminate hands in clinical situations when hands are not visibly soiled as they provide for a rapid kill of most transient microorganisms, are less time-consuming than washing with soap and water and are easier on skin.^{19, 33-36}

PERSONAL PROTECTIVE EQUIPMENT (PPE)

PPE is used to prevent transmission of infectious agents both from patient-to-patient and from patient-to-staff.

Personal protective equipment (PPE) is used alone or in combination to prevent exposure, by placing a barrier between the infectious source and one's own mucous membranes, airways, skin and clothing.^{7,18} The selection of PPE is based on the nature of the interaction with the client/patient/resident and/or the likely mode(s) of transmission of infectious agents. Selection of the appropriate PPE is based on the risk assessment (e.g., interaction, status of client/patient/resident) that dictates what is worn to break the chain of transmission. For more information about risk assessment, see Section II and [Appendix B](#).

PPE should never be used indiscriminately and overuse may have negative impacts, such as:

- a) interference with quality of client/patient/resident care^{37,38} (see also Section II, 'Impact of Isolation on Quality of Care');
- b) wastage and increased cost;
- c) staff may be less likely to wash their hands when wearing gloves for routine tasks;
- d) overuse may lead to shortages of PPE that result in inappropriate use (e.g., re-use of gloves and gowns), leading to increased transmission of microorganisms^{39,40}; and
- e) environmental concerns relating to disposable barrier equipment, washing agents and chemicals.

Personal protective equipment should be put on just prior to the interaction with the client/patient/resident. When the interaction for which the PPE was used has ended, PPE should be removed immediately and disposed of in the appropriate receptacle. The process of PPE removal requires strict adherence to a formal protocol to prevent recontamination.⁴¹ Refer to [Appendix L](#) for instructions for putting on and taking off PPE.

Health care settings must ensure that staff have sufficient supplies of, and quick, easy access to, the PPE required. Health care settings should have a process for evaluating PPE to ensure it meets quality standards where applicable,⁴ including a respiratory protection program compliant with the Ministry of Labour requirements.^{4,10}

Education in the proper use of PPE must be provided by the health care setting to all health care providers and other staff who have the potential to be exposed to blood and body fluids.

Gloves

Gloves must be worn when it is anticipated that the hands will be in contact with mucous membranes, non-intact skin, tissue, blood, body fluids, secretions, excretions, or equipment and environmental surfaces contaminated with the above.⁷

Gloves are not required for routine health care activities in which contact is limited to intact skin of the client/patient/resident (e.g., taking blood pressure, bathing and dressing the client/patient/resident). Compliance with hand hygiene should always be the first consideration.

Indiscriminate or improper glove use has been linked to transmission of pathogens.⁴² Gloves are **task-specific** and **single-use** for the task. Re-use of gloves has been associated with transmission of methicillin-resistant *Staphylococcus aureus* (MRSA) and Gram-negative bacilli.^{43,44} See [Box 2](#) for the appropriate use of gloves.

Sterile gloves are used in operating theatres and when performing sterile procedures such as central line insertions.

Selection of Gloves

It is important to assess and select the best glove for a given task. Selection of gloves should be based on a risk analysis of ⁴⁵:

- a) the type of setting (e.g., operating room, environmental cleaning, laboratory);
- b) the task that is to be performed (e.g., invasive or non-invasive);
- c) the likelihood of exposure to body substances;
- d) the anticipated length of use; and
- e) the amount of stress on the glove.

The barrier integrity of gloves varies on the basis of:

- a) type and quality of glove material;
- b) intensity of use;
- c) length of time used;
- d) manufacturer;
- e) whether gloves were tested before or after use; and
- f) method used to detect glove leaks.

It is preferable to provide more than one type of glove to health care providers, because it allows the individual to select the type that best suits their care activities¹⁸. Some additional points to consider:

- a) good quality vinyl gloves are generally sufficient for most tasks;
- b) latex or synthetic gloves, such as nitrile or neoprene gloves, are preferable for clinical procedures that require manual dexterity and/or will involve more than brief patient contact¹⁸;
- c) powdered latex gloves have been associated with latex allergy;
- d) new types of latex gloves are being developed which may be safe for those with an allergy to rubber latex⁴⁶;
- e) gloves that fit snugly around the wrist are preferred for use with a gown because they will cover the gown cuff and provide a better barrier for the arms, wrists and hands.¹⁸

- Refer to Appendix M for advantages and disadvantages of different types of medical gloves.
- For more information about standards for gloves, visit the Canadian General Standards Board website at: <http://www.pwgsc.gc.ca/cgsb/prgsrv/certprg/program/astm-5250-e.html>.

Gloves and Hand Hygiene

Because gloves are not completely free of leaks and hands may become contaminated when removing gloves,³ hands must be cleaned before putting on gloves for an aseptic/clean procedure and after glove removal.⁷ Gloves must be removed immediately and discarded into a waste receptacle after the activity for which they were used and before exiting a client/patient/resident environment.

BOX 2: Appropriate Glove Use

- **Wear the correct size of gloves.**
- **Gloves should be put on immediately before the activity for which they are indicated.**
- **Clean hands before putting on gloves for a clean/aseptic procedure.**
- **Gloves must be removed and discarded immediately after the activity for which they were used.**
- **Hand hygiene must be performed immediately after glove removal.**
- **Change or remove gloves if moving from a contaminated body site to a clean body site within the same client/patient/resident.**
- **Change or remove gloves after touching a contaminated site and before touching a clean site or the environment.**
- **Do not wash or re-use gloves.**
- **The same pair of gloves must not be used for the care of more than one client/patient/resident.**

Gloves may be adversely affected by petroleum-based hand lotions or creams. Verify with the glove manufacturer that the gloves are compatible with the hand hygiene products in use in the health care setting (e.g., lotions).

To reduce hand irritation related to gloves¹¹:

- a) wear gloves for as short a time as possible;
- b) ensure hands are clean and dry before putting on gloves; and
- c) ensure gloves are intact and clean and dry inside.

Gowns

A gown is recommended when it is anticipated that a procedure or care activity is likely to generate splashes or sprays of blood, body fluids, secretions, or excretions.⁷

Long-sleeved gowns protect the forearms and clothing of the health care provider from splashing and soiling with blood, body fluids and other potentially infectious material.

➤ See [Box 3](#) for the appropriate use of gowns.

Selection of Gowns

The type of gown selected is based on the nature of the interaction with the client/patient/resident and includes¹⁸:

- a) anticipated degree of contact with infectious material;
- b) potential for blood and body fluid penetration of the gown (e.g., water-resistant gowns should be used in the operating theatre when soaking is anticipated); and
- c) requirement for sterility (e.g., sterile gowns are worn in operating theatres and when performing sterile procedures such as central line insertions).

Gowns used as PPE should be cuffed and long-sleeved, and offer full coverage of the body front, from neck to mid-thigh or below. Clinical and laboratory coats or jackets are not a substitute for gowns where a gown is indicated. Several gown sizes should be available in a health care setting to ensure appropriate coverage for staff.

Masks

A mask is used by a health care provider (in addition to eye protection) to protect the mucous membranes of the nose and mouth when it is anticipated that a procedure or care activity is likely to generate splashes or sprays of blood, body fluids, secretions or excretions⁷ or within two metres of a coughing client/patient/resident.^{18, 47}

Masks are also required in operating theatres and when performing aseptic procedures (e.g., central line insertions, spinal epidural procedures).

BOX 3: Appropriate Gown Use

- Gowns should only be worn when providing care for clients/patients/residents.
- When use of a gown is indicated, the gown should be put on immediately before the task and must be worn properly, i.e., tied at top and around the waist.
- Remove gown immediately after the task for which it has been used in a manner that prevents contamination of clothing or skin and prevents agitation of the gown.
- Discard used gown immediately after removal into appropriate receptacle. Do not hang gowns for later use.
- Do not re-use gown. Do not go from patient-to-patient wearing the same gown.

A mask should be placed on a coughing client/patient/resident when outside their room, if tolerated, to limit dissemination of infectious respiratory secretions.¹⁸

➤ See [Box 4](#) for the appropriate use of masks.

Selection of Masks

Mask selection is based on a risk assessment that includes:

- type of procedure/care activity;
- length of procedure/care activity; and
- likelihood of contact with droplets/aerosols generated by the procedure or interaction.

Criteria for selecting masks include:

- mask should securely cover the nose and mouth;
- mask should be substantial enough to prevent droplet penetration; and
- mask should be able to perform for the duration of the activity for which the mask is indicated (e.g., surgery).

N95 Respirators

An N95 respirator is used to prevent inhalation of small particles that may contain infectious agents transmitted via the airborne route.¹⁸

N95 respirators should also be worn for aerosol-generating procedures that have been shown to expose staff to undiagnosed tuberculosis, including:

- sputum induction;
- diagnostic bronchoscopy; and
- autopsy examination.

➤ See Section II for more information about N95 respirators and their indications.

➤ Refer to [Appendix M](#) for advantages and disadvantages of different types of masks and N95 respirators.

Eye Protection

Eye protection is used by health care providers (in addition to a mask) to protect the mucous membranes of the eyes when it is anticipated that a procedure or care activity is likely to generate splashes or sprays of blood, body fluids, secretions or excretions,⁷ or within two metres of a coughing client/patient/resident.^{18, 47}

Eye protection includes:

- safety glasses;
- safety goggles;
- face shields; and
- visors attached to masks.

Prescription eye glasses are not acceptable by themselves as eye protection; they may be worn underneath face shields and some types of protective eyewear.

BOX 4: Appropriate Mask Use

- Select a mask appropriate to the activity
- Mask should securely cover the nose and mouth
- Change mask if it becomes wet.
- Do not touch mask while wearing it.
- Remove mask correctly immediately after completion of task and discard into an appropriate waste receptacle.
- Do not allow mask to hang or dangle around the neck.
- Clean hands after removing the mask.
- Do not re-use disposable masks.
- Do not fold the mask or put it in a pocket for later use.

Eye protection may be disposable or, if reusable, should be cleaned prior to re-use. Due to the risk of contamination, it is recommended that reusable eye protection be sent to a central area for reprocessing after use.

Eye protection should be comfortable, not interfere with visual acuity and fit securely. A health care setting may need to provide several different types, styles and sizes of protective eye equipment.¹⁸

➤ See [Box 5](#) for the appropriate use of eye protection.

Selection of Eye Protection

The eye protection chosen for specific situations depends on:

- the type of activity and risk of exposure;
- the circumstances of exposure (e.g., droplet exposure vs. sprays/splashes of fluid);
- other PPE used; and
- personal vision needs.

Criteria for selecting eye protection includes:

- eye protection must provide a barrier to splashes from the side;
- eye protection may be single-use disposable or washable before re-use; and
- prescription eye glasses are not acceptable as eye protection.

➤ Refer to [Appendix M](#) for advantages and disadvantages of different types of eye protection.

Routine Practices for Procedures that Generate Droplets and/or Aerosols

Certain procedures may generate droplets/aerosols that may expose staff to respiratory pathogens and are considered to be a potential risk for staff and others in the area. Personal protective equipment (mask, protective eyewear or face shield) must be used by staff when within two metres of procedures generating droplets/aerosols on any client/patient/resident, with or without symptoms of an acute respiratory infection, to prevent deposition of droplets/aerosols on staff mucous membranes.⁴⁸ See [Box 6](#) for a list of procedures that generate droplets/aerosols.

There is debate about whether other medical procedures may result in generation of droplets/aerosols and lead to transmission of respiratory infection. For these procedures, to date, there is inconclusive or no published literature documenting transmission. Examples of such procedures include:

- nebulized therapies
- high-frequency oscillatory ventilation
- performing a tracheostomy or providing tracheostomy care
- chest physiotherapy
- nasopharyngeal swabs or nasopharyngeal aspirates
- tube or needle thoracostomy.

For these procedures, use of personal protective equipment should be determined by risk assessment.

Facial protection is also required routinely for:

- breaches to the integrity of a mechanical ventilation system (e.g., open suctioning, filter changes); and
- disposal of filters used in mechanical ventilation and cleaning/disposal of bags and filters.

All units and crash carts should be equipped with:

BOX 5: Appropriate Use of Eye Protection

- Eye protection must be removed immediately after the task for which it was used and discarded into waste or placed in an appropriate receptacle for cleaning.
- Prescription eye glasses are not acceptable as eye protection.

- a) a manual resuscitation bag with hydrophobic submicron filter;
- b) in-line suction catheters for adults;
- c) non-rebreather mask that allows filtration of exhaled gases; and
- d) personal protective equipment (gloves, gowns, masks, eye protection).

BOX 6: Procedures Generating Droplets/Aerosols

- Endotracheal intubation, including during cardio-pulmonary resuscitation
- Cardio-pulmonary resuscitation
- Open airway suctioning
- Bronchoscopy*
- Surgery and autopsy
- Sputum induction*
- Non-invasive positive pressure ventilation for acute respiratory failure (CPAP, BiPAP¹⁻³)
- High concentration oxygen therapy (50% or higher)^{1, 2}

* For diagnostic (but not therapeutic) bronchoscopy or sputum induction, use an N95 respirator, due to risk from undiagnosed TB

ENVIRONMENTAL CONTROLS

Environmental controls are measures that are built into the infrastructure of the health care setting that have been shown to reduce the risk of infection to staff and clients/patients/residents, such as appropriate accommodation and placement, hand washing sinks, point-of-care alcohol-based hand rub, patient care equipment that is in good repair, cleaning practices and engineering controls such as point-of-care sharps containers and sufficient air changes per hour appropriate to the care setting. Engineering controls are the preferred controls as they do not depend on individual health care provider compliance.

Accommodation and Placement

Single rooms, with dedicated bathroom and sink, are preferred for placement of all clients/patients/residents.⁴⁹ Studies have shown a clear relationship between the use of single rooms and reduced infection.⁵⁰⁻⁵² However, most health care facilities do not have sufficient single rooms to accommodate all clients/patients/residents, so some might be accommodated in multi-bed rooms, which presents a risk for transmission of microorganisms. Clear protocols must be in place regarding patient placement in order to minimize the transmission risk to others.

In health care settings that do not have sufficient single rooms available for all routine care, decisions must be made regarding room assignments and selection of roommates based on:

- a) route of transmission of the infectious agent (known or suspected);
- b) client/patient/resident risk factors for transmission (e.g., hygiene, cognitive status);
- c) risk factors for acquisition in other clients/patients/residents in the unit (e.g., compromised immunity); and
- d) availability of single rooms.

Decision-making regarding accommodation should include the questions listed in Box 7.²⁸

BOX 7: Questions to Ask When Determining Placement of Clients/Patients/Residents and their Roommates

- Is the client/patient/resident soiling his/her environment because of poor hygiene practices, uncontained drainage or incontinence?
- Does the client/patient/resident have an infection that might be transmitted to another client/patient/resident?
- What is the condition of other clients/patients/residents in the unit?
- Does the client/patient/resident have an indwelling device (e.g., urinary catheter, central line, feeding tube)?
- Does the client/patient/resident have non-intact skin?
- What is the susceptibility level of the client/patient/resident with respect to underlying diseases, neutropenia, extremes of age?
- Is the client/patient/resident at risk for an antibiotic-resistant organism?
- Can the client/patient/resident follow directions on hygiene measures?

For clients/patients/residents who have a cough or other symptoms of an acute respiratory infection:

- a) move out of waiting area to a separate area or room (preferably with negative pressure) as soon as possible;
- b) if single room accommodation is unavailable, maintain a spatial separation of at least two metres¹⁸ between the coughing client/patient/resident and others in the room and draw the privacy curtain between beds;
- c) if there is a suspicion that the infection is transmitted via the airborne route, the client/patient/resident must be moved into a single room, preferably with negative pressure;
- d) a mask and instruction in hand hygiene and respiratory etiquette should be provided to the client/patient/resident;
- e) symptomatic clients/patients/residents should be assessed as soon as possible.

Environment and Equipment Cleaning

The physical environment of a health care setting can harbour many microorganisms that are capable of causing infection in susceptible individuals. Maintaining a clean and safe health care environment is an essential component of infection prevention and control and is integral to the safety of clients/patients/residents and staff.⁵³⁻⁵⁵

Numerous studies have shown that the inanimate health care environment harbours bacteria and viruses that may be transferred to clients/patients/residents and equipment via the hands of health care providers.^{56,57} Some studies have shown that environmental strains of microorganisms are identical to those of the client/patient/resident occupying the environmental space.^{58,59} In some instances, health care-associated infection outbreaks have been brought under control when the intensity of environmental cleaning was increased.⁶⁰

Health care settings must devote adequate resources to Environmental Services/Housekeeping that include^{9,10,18,54}:

- a) adequate human resources;
- b) availability of appropriate cleaning products;
- c) written policies and procedures for cleaning and disinfection of client/patient/resident rooms and equipment that includes cleaning standards and frequencies;
- d) education and training of cleaning staff;
- e) procedures and increased capacity for outbreak management; and

- f) ongoing review and monitoring of practices and procedures.

Policies and procedures should address the environmental aspects of areas when the role of the environment may be a significant factor in the prevention of HAIs, such as:

- a) cleaning and disinfection of non-critical equipment between clients/patients/residents, including transport equipment^{9, 54, 61};
- b) minimum high-level disinfection of semi-critical and sterilization of critical medical equipment^{12, 54};
- c) daily and terminal cleaning of rooms;
- d) cleaning requirements for rooms that house clients/patients/residents with *C.difficile* or VRE⁵⁴;
- e) management of linen and waste^{9, 54}; and
- f) cleaning in areas adjacent to construction activities^{9, 54} at the end of the day or at other times as required to maintain cleanliness.

Environmental cleaning in the health care facility should be performed on a routine and consistent basis to provide for a safe and sanitary environment.^{9, 54} Cleaning staff require education and training that includes clear messaging regarding their role in the prevention of infections in their health care setting. Cleaning practices in the health care setting must be audited and results reported back appropriately.¹⁰ Frequent audits of practice must be included as part of the organization's responsibility to maintaining a clean environment.⁹

Health care settings must review their cleaning and disinfection methods to ensure that they are adequate for disinfection of contaminated surfaces. Cleaning and disinfecting products used in the health care setting must be approved by Infection Prevention and Control and Occupational Health.¹² Hospital-grade disinfectants must have a drug identification number (DIN) from Health Canada to indicate approval for use in Canada.⁴⁵ Manufacturers' recommendations for use and dilution must be followed.^{45, 54}

For a detailed discussion regarding the implementation of a cleaning and reprocessing program, refer to:

- *Best Practices for Cleaning, Disinfection and Sterilization in All Health Care Settings*,¹² available at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/cleaning-disinfection-and-sterilization.html>
- *Best Practices for Environmental Cleaning in All Health Care Settings*,⁹ available at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/environmental-cleaning-for-prevention-and-control-of-infections.html>

Dishware and Eating Utensils¹⁸

Dishware and eating utensils are effectively decontaminated in commercial dishwashers with hot water and detergents. Reusable dishware and utensils may be used for all patients/residents including those on Additional Precautions. Disposable dishes are not required.

Food premises must comply with the requirements of the *Health Protection and Promotion Act*, R.R.O. 1990, *Regulation 562*, Food Premises,⁶² available at:

http://www.e-laws.gov.on.ca/Download?dDocName=elaws_regs_900562_e.

Linen and Waste

Laundry

Policies and procedures should address the collection, transport, handling, washing and drying of soiled linen, including protection of staff and hand hygiene. Laundry regulations should be addressed if the facility does its own laundry.⁹

- For detailed information about the management of laundry, refer to *Best Practices for Environmental Cleaning in All Health Care Settings*,⁹ available at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/environmental-cleaning-for-prevention-and-control-of-infections.html>

Linen that is soiled with blood, body fluids, secretions or excretions should be handled using the same precautions, regardless of whether the client/patient/resident is on Additional Precautions and regardless of the source or health care setting.^{9, 45, 54} In particular:

- a) bag or otherwise contain contaminated laundry at the site of collection;
- b) use leak-proof containment for laundry contaminated with blood or body substances (water soluble bags and 'double-bagging' are not recommended);
- c) laundry carts or hampers used to collect or transport soiled linen need not be covered; and
- d) linen bags should be tied securely and not over-filled.

Facilities for hand hygiene must be readily available in laundry areas and staff should clean their hands whenever gloves are changed or removed. Laundry staff should protect themselves from potential cross-infection from soiled linen by wearing appropriate protective equipment, such as gloves and gowns or aprons, when handling soiled linens.

Staff in health care areas need to be aware of sharps when placing soiled linen in bags; laundry staff are at risk from contaminated sharps, instruments or broken glass that may be contained with linen in the laundry bags. Laundry staff should be trained in procedures for safe handling of soiled linen and must be offered immunization against hepatitis B.^{9, 45}

Waste Management

Written policies and procedures for management of contaminated infectious waste from health care settings must be developed based on provincial regulations and local bylaws and should address issues such as the collection, storage, transport, handling and disposal of contaminated waste, including sharps and biomedical waste.⁹

- For more information about waste management, refer to *Best Practices for Environmental Cleaning in All Health Care Settings*, available at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/environmental-cleaning-for-prevention-and-control-of-infections.html>

Waste handlers should wear protective apparel appropriate to their risk (e.g., gloves, protective footwear). Waste handlers that may be exposed to biomedical waste and/or sharps must be offered hepatitis B immunization.

Handling of Sharps

Sharps are devices that can cause occupational injury to staff. Some examples of sharps include needles, lancets, blades and clinical glass. A sharps injury prevention program must be in place in all health care settings.^{10, 16} This should include follow-up for exposure to bloodborne pathogens.⁶³

Prevention of sharps injuries may be achieved by:

- a) the use of safety-engineered devices;
 - b) the provision of puncture-resistant sharps containers at point-of-care; and
 - c) staff education regarding the risks associated with unsafe procedures such as recapping.
- For specific requirements under Ontario's needle safety legislation see the *Occupational Health and Safety Act*, O. Regulation 474/07, Needle Safety,⁶⁴ available at: http://www.e-laws.gov.on.ca/Download?dDocName=elaws_regs_070474_e.

ADMINISTRATIVE CONTROLS

Administrative controls are measures that the health care setting puts into place to protect staff and clients/patients/residents from infection.

Staff Education and Training

Infection prevention and control education should be provided to all staff, especially those providing direct client/patient/resident care, at the initiation of employment as part of their orientation and as ongoing continuing education on a scheduled basis.¹⁰ Education in infection prevention and control must span the entire health care setting and be directed to all who work in that setting. Health care facilities should ensure that appropriate policies and procedures are in place to ensure attendance at training/education in Routine Practices and Additional Precautions (including hand hygiene) and that attendance is recorded and reported back to the manager to become a part of the employee's performance review.^{10, 65}

Effective infection prevention and control education programs should address¹⁰:

- a) disease transmission, the risks associated with infectious diseases and basic epidemiology of health care-associated infections specific to the care setting;
- b) hand hygiene, including the use of alcohol-based hand rubs and hand washing¹¹;
- c) principles and components of Routine Practices as well as Additional Precautions;
- d) assessment of the risk of exposure and the appropriate use and indications for PPE, including safe application, removal and disposal;
- e) appropriate cleaning and/or disinfection of health care equipment, supplies and surfaces or items in the health care environment^{9, 12};
- f) individual staff responsibility for keeping clients/patients/residents, themselves and co-workers safe; and
- g) education in early problem or symptom recognition.

Education of Clients/Patients/Residents

Client/patient/resident teaching should include:

- a) correct hand hygiene;
- b) basic hygiene practices that prevent the spread of microorganisms, such as respiratory etiquette; and
- c) not sharing personal items.

Client/patient/resident education about any precautions that might be required is important, as it involves them in this aspect of their care and leads to increased patient satisfaction.⁶⁶ Infection Prevention and Control may assist staff in education of clients/patients/residents through developing and/or reviewing informational materials pertaining to Routine Practices.

Respiratory Etiquette

Health care settings should reinforce with staff, clients/patients/residents and visitors the personal practices that help prevent the spread of microorganisms that cause respiratory infections. These personal practices include:

- a) not visiting people in a health care facility when acutely ill with a respiratory infection;
- b) avoidance measures that minimize contact with droplets when coughing or sneezing, such as:
 - i) turning the head away from others;
 - ii) maintaining a two-metre separation from others⁴⁷;
 - iii) covering the nose and mouth with tissue;
- c) immediate disposal of tissues into waste after use; and
- d) immediate hand hygiene after disposal of tissues.

Healthy Workplace Policies

All health care settings should establish a clear expectation that staff do not come into work when ill with symptoms that are of an infectious origin, and support this expectation with appropriate attendance management policies.⁵⁵ Staff carrying on activities in a health care setting who develop an infectious illness may be subject to some work restrictions.

The Communicable Disease Surveillance Protocols from the Ontario Hospital Association (OHA)/Ontario Medical Association (OMA)/MOHLTC state: *'Health care workers have a responsibility to their patients and colleagues regarding not working when ill with symptoms that are likely attributable to an infectious disease. This includes staff with influenza-like illness, febrile respiratory illness, gastroenteritis and conjunctivitis'*.⁶⁷

Immunization

Client/Patient/Resident Immunization

One of the most effective preventive measures to protect clients/patients/residents and staff from acquiring communicable diseases is immunization. All health care settings should have an age-appropriate immunization program in place.¹⁰

Staff Immunization

Immunization programs are highly effective and are a critical component of the Occupational Health program.^{68, 69} Health care providers must be offered appropriate immunizations. Immunizations should be based on requirements such as OHA/OMA/MOHLTC communicable disease surveillance protocols^{63, 70-72} and be consistent with recommendations from the National Advisory Committee on Immunization for health care providers.⁷³ Appropriate vaccine use protects the health care provider, colleagues and the client/patient/resident. Vaccines appropriate for health care providers include:

- a) annual Influenza vaccine⁷⁰;
- b) measles,⁷² mumps,⁷⁴ rubella⁷¹ (MMR) vaccine;
- c) varicella⁷⁵ vaccine;
- d) hepatitis B⁶³ vaccine - staff who use sharps or who may be exposed to contaminated sharps in their work should be offered hepatitis B vaccination, followed by serology to document immunity; and
- e) acellular pertussis⁷⁶ vaccine.

➤ Information regarding the Communicable Disease Surveillance Protocols is available at:

http://oha.ca/Client/OHA/OHA_LP4W_LND_WebStation.nsf/page/Communicable+Diseases+Surveillance+Protocols.

Recommendations for Routine Practices

1. ***The elements of Routine Practices must be incorporated into the culture of all health care settings and into the daily practice of each health care provider during the care of all clients/patients/residents at all times. [BII]***
2. ***Visitors should receive instruction regarding specific facility control measures before they visit a client/patient/resident, to ensure compliance with established practices. [BII]***
3. ***Perform a risk assessment before each interaction with a client/patient/resident or their environment in order to determine which interventions are required to prevent transmission during the planned interaction. [BIII]***
4. ***Choose client/patient/resident accommodation based on the risk assessment.***
5. ***Choose personal protective equipment based on the risk assessment.***
6. ***All health care settings must implement a comprehensive hand hygiene program that follows the best practices recommended in 'Best Practices for Hand Hygiene in All Health Care Settings'.***

7. *Provide sufficient supplies of easily accessible PPE. [AIII]*
8. *Implement a process for evaluating PPE to ensure it meets quality standards where applicable, including a respiratory protection program compliant with the Ministry of Labour requirements. [AIII]*
9. *Provide education in the proper use of PPE to all health care providers and other staff who have the potential to be exposed to blood and body fluids. [BII]*
10. *Wear gloves when it is anticipated that the hands will be in contact with mucous membranes, non-intact skin, tissue, blood, body fluids, secretions, excretions, or equipment and environmental surfaces contaminated with the above. [AII]*
11. *Gloves are not required for routine health care activities in which contact is limited to the intact skin of the client/patient/resident. [AIII]*
12. *Select gloves that fit well and are of sufficient durability for the task. [AII]*
13. *Put on gloves just before the task or procedure that requires them. [AII]*
14. *Perform hand hygiene before putting on gloves for aseptic procedures. [AIII]*
15. *Remove gloves immediately after completion of the task that requires gloves, before touching clean environmental surfaces. [AIII]*
16. *Clean hands immediately after removing gloves. [AII]*
17. *Single-use disposable gloves should not be re-used or washed. [AII]*
18. *Wear a gown when it is anticipated that a procedure or care activity is likely to generate splashes or sprays of blood, body fluids, secretions, or excretions. [BIII]*
19. *Remove gown immediately after the task for which it has been used in a manner that prevents contamination of clothing or skin and prevents agitation of the gown. [BII]*
20. *Wear a mask and eye protection to protect the mucous membranes of the eyes, nose and mouth when it is anticipated that a procedure or care activity is likely to generate splashes or sprays of blood, body fluids, secretions or excretions. [AII]*
21. *Wear an N95 respirator to prevent inhalation of small particles that may contain infectious agents transmitted via the airborne route. [AII]*
22. *Single rooms, with dedicated bathroom and sink, are preferred for placement of all clients/patients/residents. [BII]*
23. *If single rooms are limited, there should be clear protocols for determining options for patient placement and room sharing based on a risk assessment. [BII]*
24. *Clients/patients/residents who visibly soil the environment or for whom appropriate hygiene cannot be maintained should be placed in single rooms with dedicated toileting facilities. [AIII]*
25. *A sharps injury prevention program must be in place in all health care settings. [AII]*
26. *Appropriate policies and procedures are in place to ensure staff attendance at training/education in Routine Practices (including hand hygiene) and attendance is recorded and reported back to the manager to become a part of the employee's performance review. [AII]*
27. *There is a program that promotes respiratory etiquette to staff, clients/patients/residents and visitors in the health care setting. [AII]*
28. *There is a clear expectation that staff do not come into work when ill with symptoms that are of an infectious origin, and this expectation is supported with appropriate attendance management policies. [BII]*

Additional Precautions

Additional Precautions refer to infection prevention and control interventions (e.g., barrier equipment, accommodation, additional environmental controls) to be used in addition to Routine Practices to protect staff and clients/patients/residents to interrupt transmission of infectious agents that are suspected or identified in a client/patient/resident.

- Refer to Appendix N for infectious diseases and agents that require Additional Precautions.

Additional Precautions are based on the mode of transmission (e.g., direct or indirect contact, airborne or droplet). There are three categories of Additional Precautions: Contact Precautions, Droplet Precautions and Airborne Precautions.

ELEMENTS THAT COMPRISE ADDITIONAL PRECAUTIONS

In addition to Routine Practices, the following elements comprise Additional Precautions:

Specialized Accommodation and Signage

Specialized accommodation and signage for clients/patients/residents on Additional Precautions includes:

- spatial separation**, such as single room⁵² and private toileting facilities for clients/patients/residents on Additional Precautions:
 - in some cases where clients/patients/residents are known to be infected with the same microorganism, cohorting is acceptable;
 - refer to Appendix C for accommodation recommendations.
- signage** specific to the type(s) of Additional Precautions:
 - a sign that lists the required precautions should be posted at the entrance to the client/patient/resident's room or bed space;
 - signage should maintain privacy by indicating only the precautions that are required, not information regarding the patient's condition;
 - refer to Appendixes F-K for sample signage.
- specialized **engineering controls** may be required for some types of Additional Precautions, e.g., negative pressure ventilation for Airborne Precautions. See Section II for information regarding engineering controls for airborne isolation rooms.

Barrier Equipment

Personal protective equipment (i.e., barrier equipment) is standardized and specific to the type(s) of Additional Precautions that are in place, e.g., gloves are required for entry to a Contact Precautions room regardless of the interactions that are to take place. If the health care provider needs to leave the room, the PPE must be removed and discarded. Fresh PPE must be worn if the health care provider re-enters the room.

Dedicated Equipment

Equipment must be dedicated to the client/patient/resident whenever possible. Equipment and supplies that are required for the interaction should be assembled first and brought into the room after PPE has been put on.

Additional Cleaning Measures

Additional cleaning measures may be required for the client/patient/resident environment. The need for cleaning some items might be reduced if they are covered with a disposable or washable sheet before use (e.g., wheelchair, couch).

- For more information about environmental cleaning in health care settings, refer to the *Best Practices for Environmental Cleaning in All Health Care Settings*⁹, available at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/environmental-cleaning-for-prevention-and-control-of-infections.html>

Limited Transport Procedures


Transport of clients/patients/residents on Additional Precautions may be limited in some cases. The following points must be considered:

- normal health care activities must be maintained despite Additional Precautions, to ensure quality of care (e.g., ambulation as part of recovery from hip surgery);
- clients/patients/residents who leave their room must be assessed to determine their risk of transmission to others; and
- for some conditions, limit transport of the client/patient/resident unless medically necessary (e.g., tuberculosis, acute viral respiratory illness, acute viral gastroenteritis such as Norovirus infection).

Communication

Effective communication regarding Additional Precautions is essential when a client/patient/resident goes to another department for testing, to another unit or to other health care settings/facilities. This communication must include Emergency Medical Services (EMS) staff and other transport staff.

- See [Box 8](#) for a summary of the elements that comprise Additional Precautions.



COHORTING

Cohorting practices can be utilized when single rooms are not available or during outbreak situations. Cohorting has contributed to the control of a number of outbreaks^{77, 78} and should be considered when transmission is documented and continues despite alternative interventions, and when available facilities and staffing allow for the establishment of cohorting.⁷⁹ Cohorting should never compromise infection control practices and Additional Precautions must be applied individually for each patient/resident within the cohort.

Cohorting consists of two components: patient/resident cohorting and staff cohorting. Patient cohorting is applicable in acute care settings for control of transmission of microorganisms or outbreaks. In long-term care homes, movement of residents to achieve geographical cohorting is not appropriate because displacement of residents from their own rooms will often cause harm to the resident, e.g., anxiety, disorientation. Staff cohorting is applicable in all health care facilities.

Patient/Resident Cohorting

Patient/resident cohorting refers to:

- a) the placement and care of individuals who are infected or colonized with the same microorganism in the same room; or
- b) placing those who have been exposed together to limit risk of further transmission.

In long-term care homes, resident cohorting does not imply that a resident is moved out of his/her room. Rather, those who already share a room or who share a bathroom and who are infected or colonized with the same microorganism may be treated as a cohort.

Geographical cohorting refers to restricting patients who are infected or colonized with the same microorganism to several rooms along a corridor or an entire clinical unit. Use of this practice can limit transmission by segregating those who are infected or colonized to a specified area away from those who are not.⁸⁰

Care equipment must be dedicated or cleaned between use on patients/residents in the same room and protective barriers such as gowns and gloves should be worn for the care of an individual patient/resident only and not worn from patient-to-patient within the cohort.

Care should be taken to assess patients/residents for the duration of colonization/infection. Avoid placement of newly identified cases together with those who have a longer history of acquisition (who may no longer be infected or colonized with the microorganism) to prevent re-exposure.

Staff Cohorting

Staff cohorting is the practice of assigning specified health care providers to care only for clients/patients/residents known to be colonized or infected with the same microorganism. These health care providers would not participate in the care of clients/patients/residents who are not colonized or infected with that microorganism.

Staff cohorting can be used in addition to client/patient/resident and geographical cohorting by assigning dedicated staff to care for either those patients/residents who are infected or colonized, or those who are not. This practice can be used during outbreaks to reduce the potential for cross-infection between clients/patients/residents by limiting the number of staff interacting with clients/patients/residents.⁸¹⁻⁸³ It can also be used to limit the number of health care providers exposed to infected cases.⁸¹

ADDITIONAL PRECAUTIONS FOR VISITORS

Visitors of clients/patients/residents on Additional Precautions in health care facilities:

- a) should be kept to a minimum;
- b) must receive education regarding hand hygiene and the appropriate use of PPE as described under Routine Practices; and
- c) must wear the same personal protective equipment as health care providers if in contact with other clients/patients/residents or providing direct care.

Clients/patients/residents and visitors must be informed about the reason for implementing Additional Precautions and receive instruction regarding how to enter and leave the room safely when the client/patient/resident is on Additional Precautions. This should include demonstration in putting on, taking off and disposing of PPE as required, as well as hand hygiene.

INITIATION AND DISCONTINUATION OF ADDITIONAL PRECAUTIONS

When Additional Precautions are instituted, they are always used in addition to Routine Practices.

Initiation of Additional Precautions

Additional Precautions must be instituted as soon as symptoms suggestive of an infection are noted, not only when a diagnosis is confirmed (see [Box 9](#) for examples). Instituting Additional Precautions should be considered before laboratory confirmation of status for patients believed to be at particularly high risk of being colonized or infected with antibiotic-resistant organisms (AROs) such as MRSA or VRE, in accordance with the health care setting's policy.

Each health care setting should have a policy authorizing any regulated health care professional to initiate the appropriate Additional Precautions at the onset of symptoms and maintain precautions until laboratory results are available to confirm or rule out the diagnosis. The person designated as the Infection Control Professional (ICP) for the health care setting:

- a) must be informed when Additional Precautions are initiated;
- b) will verify that the precautions are appropriate to the situation; and
- c) will be consulted before discontinuation of Additional Precautions or as per health care setting policy.

BOX 9: Clinical Syndromes Requiring the Use of Controls (including PPE) Pending Diagnosis

- **Acute diarrhea and/or vomiting of suspected infectious etiology:**
 - GLOVES, SINGLE ROOM
 - GOWN if skin or clothing will come into direct contact with the patient or the patient's environment and for paediatrics and incontinent/non-compliant adults
- **Acute respiratory infection, undiagnosed:**
 - SINGLE ROOM/SPATIAL SEPARATION preferred, FACIAL PROTECTION, GLOVES
 - GOWN if skin or clothing will come into direct contact with the patient or the patient's environment
- **Respiratory infection with risk factors and symptoms suggestive of Tuberculosis:**
 - FIT-TESTED N95 RESPIRATOR, NEGATIVE PRESSURE ROOM
- **Suspected meningitis and/or sepsis with petechial rash:**
 - SINGLE ROOM, FACIAL PROTECTION
- **Undiagnosed rash without fever:**
 - GLOVES
- **Rash suggestive of varicella or measles:**
 - NEGATIVE PRESSURE ROOM – only immune staff to enter
- **Abscess or draining wound that cannot be contained:**
 - GLOVES
 - GOWN if skin or clothing will come into direct contact with the patient

Duration and Discontinuation of Additional Precautions

The health care setting should have a policy that permits **discontinuation of Additional Precautions in consultation with the Infection Prevention and Control Professional** or designate. The attending physician should be notified when Additional Precautions are being discontinued. If there is disagreement between the ICP and the attending physician regarding the discontinuation, then the higher level of precautions will remain in effect with daily review until there is a definitive diagnosis or expert consultation.

Additional Precautions should remain in place until there is no longer a risk of transmission of the microorganism or illness. In some instances expert consultation may be required.

Health care settings should have policies that authorize the Infection Prevention and Control Professional to initiate and/or discontinue Additional Precautions.

Where the periods of communicability are known, precautions may be discontinued at the appropriate time.

- Refer to [Appendix N](#), 'Clinical Syndromes and Conditions with Required Level of Precautions', for recommendations related to the duration of Additional Precautions for specific illnesses.

For recommendations for discontinuation of precautions for methicillin-resistant *Staphylococcus aureus* (MRSA), VRE and *C. difficile*:

- For MRSA and VRE, refer to *Annex A: Screening, Testing and Surveillance for Antibiotic-Resistant Organisms (AROs) in All Health Care Settings*, available on the PHO website at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/screening-testing-and-surveillance-for-antibiotic-resistant-organisms-aros.html>
- For *C.difficile*, refer to *Annex C: Testing, Surveillance and Management of Clostridium difficile in All Health Care Settings*, available on the PHO website at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/testing-surveillance-and-management-of-clostridium-difficile.html>

Impact of Additional Precautions on Quality of Care^{37, 38, 84-86}

Although Additional Precautions, such as wearing gloves and single room accommodation, are necessary to protect both other clients/patients/residents and health care providers, there are negative impacts for the client/patient/resident. These include⁸⁵:

- limited contact with health care providers may result in lack of monitoring processes such as recording of vital signs and physician visits,⁸⁶ medication errors, increases in falls;
- fewer visits from family and friends, often resulting in feelings of loneliness and interfering in needed emotional support; and
- psychological problems related to isolation such as anxiety, depression, sleep disturbance, withdrawal, regression and hallucinations.

Psychological support for the client/patient/resident may include structured recreation programs, steps to prevent time disorientation and psychological support for both clients/patients/residents and their families.⁸⁵

It is important that Additional Precautions not be used any longer than necessary and that frequent assessment of the risks of transmission be carried out by infection prevention and control professionals with the goal being the removal of precautions as soon as it is safe to do so. Modification of precautions may be required for medical purposes (e.g., to permit specialized testing) or on compassionate grounds.

CONTACT TRANSMISSION AND CONTACT PRECAUTIONS

Contact Precautions are used in addition to Routine Practices for microorganisms where contamination of the environment or intact skin is a particular consideration, such as:

- a) contamination of the client/patient/resident environment;
- b) infectious agents of very low infective dose (e.g., Norovirus, rotavirus); and
- c) clients/patients/residents infected or colonized with epidemiologically important microorganisms that may be transmitted by contact with intact skin or with contaminated environmental surfaces (e.g., MRSA, VRE, *C. difficile*).⁷

Contact Transmission

Contact transmission is the most common route of transmission of infectious agents. There are two types of contact transmission:

- a) *Direct contact* occurs through touching; for example, an individual may transmit microorganisms to others by touching them; and
- b) *Indirect contact* occurs when microorganisms are transferred via contaminated objects or the hands of a health care provider coming into contact with an individual; for example, *C. difficile* might be transferred between patients, if a commode used by a patient with *C. difficile* is taken to another patient without cleaning and disinfecting the commode in between uses.

Microorganisms transmitted by contact transmission include many of the epidemiologically significant infections in health care settings: methicillin-resistant *Staphylococcus aureus* (MRSA), vancomycin-resistant enterococcus (VRE), *Clostridium difficile*, *Acinetobacter baumannii* and the agents of infectious diarrheas.

Rationale for Barrier Equipment to Reduce Contact Transmission

Several studies provide evidence that the appropriate use of gloves can help reduce transmission of pathogens in health care settings.⁸⁷⁻⁸⁹ Gown use has been shown to be effective in the control of epidemiologically important pathogens, such as VRE.^{56, 90-93}

Elements that Comprise Contact Precautions

In addition to Routine Practices, the elements that comprise Contact Precautions are summarized in [Table 2](#).

Contact Precautions are always in addition to Routine Practices such as hand hygiene. Ensure **hand hygiene by the patient** on leaving his/her room. Clients/patients/residents should be encouraged to perform hand hygiene on presentation and departure from an ambulatory/clinic setting.

Accommodation

Preferred accommodation in acute care for Contact Precautions is a single room with a dedicated toilet and patient sink.^{7, 18} The door may remain open. If single rooms are unavailable, clients/patients/residents may be cohorted with other clients/patients/residents who are infected with the same microorganism.

In long-term care and other residential settings, placement of residents requiring Contact Precautions should be reviewed on a case-by-case basis.¹⁸ Infection risk to other occupants of the room must be considered when selecting roommates.

In ambulatory settings, place patients who require Contact Precautions in an examination room or cubicle as soon as possible.¹⁸

- [Refer to Appendix C](#), 'Decision-Making Related to Accommodation and Additional Precautions', for a guide to assist with the accommodation and placement of clients/patients/residents requiring Contact Precautions.

Cleaning and Transport

Routine cleaning practices are acceptable for most rooms on Additional Precautions. Modified or additional environmental cleaning procedures and transportation of clients/patients/residents with AROs are important components of Contact Precautions for VRE and *C. difficile*.

For more information:

- Refer to *Annex A: Screening, Testing and Surveillance for Antibiotic-Resistant Organisms (AROs) in All Health Care Settings*, available on the PHO website at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/screening-testing-and-surveillance-for-antibiotic-resistant-organisms-aros.html>
- Refer to *Annex C: Testing, Surveillance and Management of Clostridium difficile in All Health Care Settings*, available on the PHO website at: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/testing-surveillance-and-management-of-clostridium-difficile.html>

Visitors

Visitors should receive education regarding hand hygiene. PPE is not required unless the visitor is providing direct care.

Barrier Equipment/PPE

- a) In acute care, gloves must be worn on entering the patient's room or bed space. Gloves must be removed and hands cleaned on exit from the room or bed space.
- b) In acute care, a gown must be worn if skin or clothing will come in contact with the patient or the patient's environment. For example:
 - a gown is required:
 - in rooms of children who are incontinent or too immature to comply with hygiene
 - in rooms of non-compliant adults who soil the environment
 - in crowded rooms/bed spaces where there is a likelihood of coming into contact with contaminated furnishings or equipment
 - when providing direct care, such as physical examination, checking vital signs, bathing or turning the patient, changing clothing, continence care, dressing changes, care of open wounds
 - a gown is not required:
 - when delivering a food tray
 - when doing a visual check of a patient at night
 - when speaking to a patient without touching anything

If a health care provider enters a Contact Precautions room without a gown and is then required to perform an activity that requires a gown, he/she must exit the room, put on a gown, then return to the room. If worn, the gown must be removed and hands cleaned on exit from the room.

- c) In non-acute settings, gloves and gown are required for activities that involve direct care (see glossary) where the health care provider's skin or clothing may come in direct contact with the resident or items in the resident's room or bed space. Gloves and gown, if worn, must be removed and hands cleaned immediately following the activity for which they were used.
- d) It is never appropriate for clients/patients/residents to wear gloves or isolation gowns while outside their room.

TABLE 2: ELEMENTS THAT COMPRISE CONTACT PRECAUTIONS

NOTE: Interventions listed in this table are in addition to Routine Practices

Element	Acute Care	Complex Continuing Care/Rehab	Long-term Care	Ambulatory/ Clinic Setting	Home Health Care
Accommodation	Door may be open				No restrictions on accommodation
	Single room with dedicated toilet and patient sink		Placement is on a case-by-case basis		
	Remain in room unless required for diagnostic, therapeutic or ambulation purposes	Not required to remain in room unless symptomatic		Identify patients who require precautions	
	May go, or be taken, outside the facility, but cannot visit other patient rooms			Encourage client to perform hand hygiene on entering the setting	
Signage	Yes			Flag chart	
Gloves	For all activities in the room/bed space		For direct care (see glossary)		
Gown	For all activities where skin or clothing will come in contact with the patient or the patient's environment		For direct care (see glossary)		
Equipment and items in the environment	Dedicate if possible			As per Routine Practices	As per Routine Practices
	Chart (paper or mobile electronic) should not be taken into the room	Clean and disinfect shared items (e.g., assigned dining area) or cover with a sheet before use		Clean and disinfect shared items (e.g., chair, examination table) or cover with a sheet before use	
Environmental Cleaning	VRE and <i>C. difficile</i> rooms require special cleaning				No special cleaning requirements
	Routine cleaning for all other rooms				
	Remove and launder all curtains (privacy, window, shower) when visibly soiled and on terminal cleaning				
Transport	Staff wear gloves and gown for direct contact with the patient during transport		Staff wear appropriate PPE for direct contact with the resident during transport	Not applicable	
	Clean and disinfect equipment used for transport after use				
Communication	Effective communication regarding precautions must be given to client/patient/resident, families, other departments, other facilities and transport services prior to transfer				

DROPLET TRANSMISSION AND DROPLET PRECAUTIONS

Droplet Precautions are used in addition to Routine Practices for clients/patients/residents known or suspected of having an infection that can be transmitted by large respiratory droplets.

Droplet Transmission

Droplet transmission occurs when droplets carrying an infectious agent exit the respiratory tract of a person. Droplets can be generated when he or she talks, coughs or sneezes and through some procedures performed on the respiratory tract (e.g., suctioning, bronchoscopy or nebulized therapies). These droplets are propelled a short distance and may enter the host's eyes, nose or mouth or fall onto surfaces. For example, if a person is coughed on by someone who has an acute respiratory infection and the secretions come in contact with mucous membranes, infection may be transmitted. Recent work suggests that droplets forcibly expelled from a cough or sneeze travel for up to two metres.⁹⁴ For patients who cannot cough forcibly, the distance that droplets travel will be less, e.g., infants and frail elderly.

Droplets do not remain suspended in the air and usually travel less than two metres (see [Figure 5](#)).⁹⁴ Microorganisms contained in these droplets are then deposited on surfaces in the client/patient/resident's immediate environment and some microorganisms remain viable for extended periods of time. Contact transmission can then occur by touching surfaces and objects contaminated with respiratory droplets.⁷

Microorganisms transmitted by this route are of special concern in certain populations, e.g., paediatrics, frail elderly, persons with cardiopulmonary disease.⁷ Examples of microorganisms transmitted by droplet transmission include: respiratory tract viruses (e.g., adenovirus, influenza and parainfluenza viruses, rhinovirus, human metapneumovirus, respiratory syncytial virus - RSV), rubella, mumps and *Bordetella pertussis*.



Figure 5: Droplet Transmission from Coughing or Sneezing

Elements that Comprise Droplet Precautions

In addition to Routine Practices, the elements that comprise Droplet Precautions are summarized in [Table 3](#).

Droplet Precautions are always in addition to Routine Practices such as hand hygiene. Ensure **hand hygiene by the patient** on leaving his/her room. Clients/patients/residents must perform hand hygiene on presentation and departure from an ambulatory/clinic setting.

Accommodation

Preferred accommodation for Droplet Precautions in acute care is a single room with a dedicated toilet and patient sink, and door may remain open. In long-term care, residents should remain in their room/bed space, if feasible, with privacy curtains drawn.

- Refer to Appendix C, 'Decision-Making Related to Accommodation and Additional Precautions', for a guide to assist with the accommodation and placement of clients/patients/residents who require Droplet Precautions.

TABLE 3: ELEMENTS THAT COMPRISE DROPLET PRECAUTIONS

NOTE: Interventions listed in this table are in addition to Routine Practices

Element	Acute Care	Complex Continuing Care/Rehab	Long-term Care	Ambulatory/ Clinic Setting	Home Health Care
Accommodation	Door may be open				Discuss feasibility of spatial separation with client (e.g., when sleeping)
	Single room with dedicated toilet and patient sink preferred	Patient/resident to remain in room or bed space if feasible, or wear a mask (if tolerated) if coughing within two metres of other patients, until no longer infectious		Triage client/patient away from waiting area to a single room as soon as possible, or maintain a two-metre spatial separation	
	Cohorting of those who are confirmed to have the same infectious agent may be acceptable	Draw privacy curtain		Patient to wear a mask and perform hand hygiene	
	Remain in room unless required for diagnostic, therapeutic or ambulation purposes				
Signage	Yes				Not applicable
Facial Protection	Yes, within two metres of client/patient/resident				
Equipment and items in the environment	Dedicate if possible				
	Chart (paper or mobile electronic) should not be taken into the room				
Environmental Cleaning	Routine cleaning				
Transport	Patient to wear a mask during transport Limit transport unless required for diagnostic or therapeutic procedures		Resident to wear a mask during transport	Client/patient to wear a mask for duration of visit and during transport	Not applicable
Communication	Effective communication regarding precautions must be given to patient families, other departments, other facilities and transport services prior to transfer				

Transport

In most cases, transport should be limited unless required for diagnostic or therapeutic procedures, such as ambulation. The client/patient/resident must wear a mask during transport, if tolerated. If the client/patient/resident cannot tolerate wearing a mask, transport staff should wear a mask and eye protection.

Barrier Equipment/PPE

A mask and eye protection must be worn by any individual who is within two metres of the client/patient/resident on Droplet Precautions.

Visitors

Visitors should receive education regarding hand hygiene. A mask should be worn by visitors within two meters of the client/patient/resident. For paediatrics, household contacts of children on Droplet Precautions do not need to wear PPE, as they will have already been exposed in the household.

AIRBORNE TRANSMISSION AND AIRBORNE PRECAUTIONS

Airborne Precautions are used in addition to Routine Practices for clients/patients/residents known or suspected of having an illness transmitted by the airborne route (i.e., particles that remain suspended in the air and may be inhaled by others).⁷

Airborne Transmission

Airborne transmission occurs when airborne particles remain suspended in the air, travel on air currents and are then inhaled by others who are nearby or who may be some distance away from the source patient, in a different room or ward (depending on air currents) or in the same room that a patient has left, if there have been insufficient air exchanges.⁷ Control of airborne transmission requires control of air flow through special ventilation systems and the use of respirators.⁷ Microorganisms transmitted by the airborne route are *Mycobacterium tuberculosis* (TB), varicella virus (chickenpox virus) and measles virus.

Effective control of airborne microorganisms hinges on maintaining a high degree of suspicion for those who present with compatible symptoms of an airborne infection,⁹⁵ early isolation in an appropriate environment and rapid diagnosis. For measles and varicella, immunization is the primary means of control.

Controls for preventing the transmission of airborne infections include:

- a) immunity against measles and varicella (immunization, natural immunity);
- b) early identification of potential cases;
- c) prompt isolation in negative-pressure airborne isolation room;
- d) appropriate treatment of client/patient/resident, where applicable;
- e) the use of a fit-tested, seal-checked N95 respirator when indicated; and
- f) identification and follow-up of exposed clients/patients/residents and staff.

Elements that Comprise Airborne Precautions

N95 Respirators

An N95 respirator must be worn when entering the room, transporting⁹⁶ or caring for a client/patient/resident with signs and symptoms or a diagnosis of active pulmonary or laryngeal tuberculosis. An N95 respirator must also be worn if non-immune staff are required to enter the room of a client/patient/resident with measles or varicella when there are no qualified immune staff available and patient safety would be compromised if they did not provide care.

N95 respirators must⁷:

- a) filter particles one micron in size;
- b) have a 95% filter efficiency; and
- c) provide a tight facial seal with less than 10% leak.

See [Box 10](#) for the appropriate use of N95 respirators.

Health care settings that use respirators must have a respiratory protection program in place. See Section II for more information on respiratory protection programs.

In health care settings specializing in care for patients with active tuberculosis (e.g., TB hospitals or units), staff may choose to re-use N95 respirators. If re-using a respirator it must be stored in a way that keeps it clean, dry, not crushed or folded and not used by anyone else. If the N95 respirator was used for a client/patient/resident who is also on Droplet or Contact Precautions, it must be discarded on removal and not re-used.

Client/Patient/Resident Controls

Patients on Airborne Precautions should remain in the airborne isolation room unless required to leave for medical reasons.

A mask is effective in trapping the large infectious particles expelled by coughing patients. Clients/patients/residents suspected or confirmed to have an airborne infection are to wear a mask at all times, if tolerated, when they must leave an area that has correct engineering controls (i.e., negative pressure ventilation). If the patient is ventilated, a filter must be present on the expiratory circuit. **There is never an indication for a client/patient/resident to wear an N95 respirator.**

Visitors

Visitors should receive education about hand hygiene. For TB, household contacts should be assessed for active tuberculosis prior to visiting the facility. Household contacts are not required to wear an N95 respirator when visiting, as they will already have been exposed in the household.

Visitors other than household contacts should be kept to a minimum and, if visiting, should be counselled and wear an N95 respirator.

Specialized Accommodation for Airborne Precautions

For clients/patients/residents on Airborne Precautions, single room accommodation in an airborne isolation room that has engineering controls in place consistent with standards from the Canadian Standards Association (CSA) is required. If an airborne isolation room is not available, transfer the patient to a facility with appropriate accommodation as soon as medically feasible. See below for engineering controls required for airborne isolation rooms.

Recommended Engineering Controls for Reducing Transmission of Microorganisms Spread by the Airborne Route

Engineering controls (e.g., directional negative pressure ventilation) are the most preferred and most effective method of minimizing exposure to airborne infections and should be used in high risk areas. Airborne isolation rooms must meet ventilation standards established by the Canadian Standards Association⁹⁷ and should meet

BOX 10: Appropriate Use of N95 Respirators

- **Select respirator for which you have been fit-tested.**
- **Perform a seal-check each time a respirator is applied.**
- **Change respirator if wet or soiled.**
- **Remove the respirator correctly and discard on removal into an appropriate receptacle.**
- **Perform hand hygiene after removing the respirator.**
- **NEVER put an N95 respirator on a client/patient/resident.**

the patient placement guidelines published by the Public Health Agency of Canada (PHAC)^{7,96} (see [Boxes 11 and 12](#) for requirements).

At a minimum, the emergency room, bronchoscopy suites, critical care settings and autopsy suites must have rooms with negative pressure capabilities as described above for high risk procedures. In acute settings expected to care for patients with infectious pulmonary tuberculosis, measles, varicella or disseminated zoster, a sufficient number of negative pressure rooms must be available on in-patient units.

An assessment of the risk of exposure to airborne infections will assist in establishing the location and number of negative pressure/ airborne isolation rooms required in order to decrease the risk of exposure to airborne infections in health care settings as described above for high risk procedures.

If using a portable HEPA-filtration unit, the ventilation requirements for an airborne isolation room as listed in [Box 11](#) must be met.

In addition to Routine Practices, the elements that comprise Airborne Precautions are summarized in [Table 4](#).^{7,18}

BOX 11: CSA Standards for Ventilation in Airborne Isolation Rooms⁹⁶

Airborne isolation rooms shall have:

- ventilation creating inward directional airflow from adjacent spaces to the room (*'negative pressure'*):
 - monitor room on initiation of use;
 - monitor at least daily when in-use;
 - monitor monthly between uses;
- an alarm indicating that the pressure relationship is not being maintained, provided just outside the room and at the nurse's station or point of supervision;
- directional airflow within the room such that clean supply air flows first to parts of the room where staff or visitors are likely to be present, and then flows across the bed area to the exhaust;
- nonaspirating diffusers;
- low-level exhaust near the head of the bed;
- exhaust air to the outdoors via dedicated exhaust:
 - washroom shall be exhausted using the same exhaust system as the room;
 - exhaust fan shall be supplied by emergency power;
- HEPA filtration of exhaust in cases where exhaust air is not discharged clear of building openings or where a risk of recirculation exists;
- minimum 12 air changes per hour;
- minimum 3 outdoor air changes per hour; and
- frequent monitoring of supply and exhaust system function by staff trained in appropriate assessment of the airflow; direction of air flow should be tested with smoke tubes at all four corners of the door.

In this Standard, the word 'shall' indicates a mandatory requirement which is the same as the use of 'shall' in this document.

BOX 12: PHAC Guidelines for Use of Airborne Isolation Rooms^{4,99}

- In acute and long-term care settings the client/patient/resident is to be placed in an airborne isolation room that meets the criteria set out by the Canadian Standards Association (see [Box 11](#), above).
- Room should have toilet, hand washing sink and bathing facilities.
- Door must be kept closed whether or not client/patient/resident is in the room.
- Windows must remain closed at all times; opening the window may cause reversal of air flow, an effect that can vary according to wind direction and indoor/outdoor temperature differentials.
- Room door must remain closed and negative airflow maintained after client/patient/resident discharge until all air in the room has been replaced; this will vary based on the number of room air changes per hour; consult facility plant engineers to determine the air changes per hour for each airborne isolation room (refer to Appendix D, 'Time Required for Airborne Isolation Room to Clear *M.tuberculosis*').
- A preventative maintenance program must be in place.
- If a **long-term care setting** does not have the appropriate facilities for airborne precautions, the resident is to be transferred to a health care facility equipped to manage airborne infections; if the transfer is delayed or not possible, place the resident in a single room with the door and window closed.
- In **ambulatory settings**, clients with suspected airborne infection should not wait in a common area but be placed directly into an examining room. Preferably this should be a negative pressure room with exhaust vented to the outside or filtered through a high efficiency filter if recirculated. If a well ventilated room is not available, a single room should be used and the client examined and discharged as quickly as possible. The door must be closed.
- In **aerosol-generating procedure rooms** where patients with airborne infections are expected to be seen (e.g., bronchoscopy suite, autopsy suite, rooms used for sputum inductions)⁹⁹:
 - there is to be a **minimum** of 12 air changes per hour in new facilities and a minimum of six air changes per hour in existing facilities;
 - the room must have inward directional air flow;
 - the air is to be exhausted directly outside the building and away from intake ducts or through a high efficiency particulate air (HEPA) filter, if recycled; and
 - the Canadian Tuberculosis Standards recommend a minimum of 15 air changes per hour for these rooms.

Adapted from Health Canada's 'Routine Practices and Additional Precautions for Preventing the Transmission of Infection in Health Care', 1999 [under revision]⁴ and the Public Health Agency of Canada's 'Canadian Tuberculosis Standards', 2007]⁹⁹

TABLE 4: ELEMENTS THAT COMPRISE AIRBORNE PRECAUTIONSNOTE: Interventions listed in this table are in addition to Routine Practices

Element	Acute Care	Complex Continuing Care/Rehab	Long-term Care	Ambulatory/ Clinic Setting	Home Health Care
Accommodation	Airborne isolation room or transfer			Airborne isolation room if available or alternate arrangements if possible	Not applicable
Signage	Yes				Not applicable
N95 Respirator	For entry to room			For duration of visit	For entry to client's home
TB	Only immune staff to enter room. N95 respirator not required if immune.				
Measles, Varicella					
Equipment and items in the environment	As per Routine Practices				
Environmental Cleaning	Routine cleaning				Routine household cleaning
Transport	Client/patient/resident to wear a mask during transport				Not applicable
	Transport staff to wear an N95 respirator during transport				
	Limit transport unless required for diagnostic or therapeutic procedures				
Communication	Effective communication regarding precautions must be given to patient families, other departments, other facilities and transport services prior to transfer				

COMBINATIONS OF ADDITIONAL PRECAUTIONS

Where more than one mode of transmission exists for a particular microorganism, the precautions used must take into consideration both modes.

Most infectious agents have a primary mode of transmission but may also have a secondary mode of transmission. Where more than one mode of transmission exists for a particular microorganism, the precautions used must take into consideration both modes. For example, respiratory viruses may remain viable for some time in droplets that have settled on objects in the immediate environment of the client/patient/resident and may be picked up on the hands of patients or staff. These microorganisms may be transmitted by contact as well as by droplet transmission and, therefore, both Contact and Droplet Precautions are required.^{7, 18}

If both tuberculosis and a respiratory virus are suspected in a single individual, a combination of Airborne, Droplet and Contact Precautions should be used. In this case, the N95 respirator must be discarded after each use and not re-used, as the outside of the respirator will be contaminated.

PROTECTIVE ENVIRONMENT

There is insufficient evidence to support the use of a protective environment (formerly known as ‘reverse isolation’) for severely immunocompromised patients such as allogeneic haematopoietic stem cell transplant (HSCT) patients and febrile neutropenic patients. These patients should be accommodated in a single room. Health care providers and others who are acutely ill with an infection should not enter the room of these patients.

Guidelines for protective environments are available from the U.S.A.:

- Healthcare Infection Control Practices Advisory Committee’s ‘*Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings 2007*’,¹⁸ available at: <http://www.cdc.gov/ncidod/dhqp/pdf/guidelines/Isolation2007.pdf>.

Recommendations for Additional Precautions

- 29. The elements of Additional Precautions must be incorporated into the health care practices of each health care setting. [BII]**
- 30. Appropriate policies and procedures are in place to ensure staff attendance at training/education in Additional Precautions and attendance is recorded and reported back to the manager to become a part of the employee’s performance review. [AII]**
- 31. When single patient rooms are limited, determine the feasibility of cohorting patients who are infected or colonized with the same microorganism. [BIII]**
- 32. Consider the use of geographic cohorting patients and staff to reduce transmission during outbreaks. [AII]**
- 33. When cohorting, Additional Precautions must be applied individually for each patient/resident within the cohort. Gowns and gloves must not be worn from patient-to-patient within the cohort and patient care equipment must not be shared.**
- 34. Visitors to clients/patients/residents on Additional Precautions must wear the same personal protective equipment as health care providers if they will be in contact with clients/patients/residents or are providing direct care. [BIII]**
- 35. Each health care setting should have a policy authorizing any regulated health care professional to initiate the appropriate Additional Precautions at the onset of symptoms. [BII]**
- 36. Additional Precautions should remain in place until there is no longer a risk of transmission of the microorganism or illness. [AII]**
- 37. The health care setting should have a policy that permits discontinuation of Additional Precautions in consultation with the Infection Prevention and Control Professional or designate. [BIII]**
- 38. Additional Precautions should not be used any longer than necessary; ongoing assessment of the risk of transmission should be performed by Infection Prevention and Control Professionals. [AII]**
- 39. In acute care, place patients who require Contact Precautions in a single room with dedicated toilet and patient sink when available. [AII]**
- 40. In long-term care and other residential settings, placement of residents who require Contact Precautions should be determined on a case-by-case basis using a risk assessment. [BII]**
- 41. In ambulatory settings, place patients who require Contact Precautions in an examination room or cubicle as soon as possible. [BII]**
- 42. In acute care, wear gloves for all activities in the patient’s room or bed space. Remove gloves and perform hand hygiene immediately on leaving the room or bed space. [AII]**
- 43. In acute care, wear a gown for all activities where skin or clothing will come in contact with the patient or the patient’s environment. When indicated, put on gown on entry to the patient’s room or bed space. If used, remove gown and perform hand hygiene immediately on leaving the room or bed space. [BIII]**
- 44. In non-acute settings, wear gloves and a gown for activities that involve direct care. Remove gloves and gown, if worn, and perform hand hygiene immediately on leaving the room. [AII]**

45. *Whenever possible, dedicate equipment and items to the patient/resident. [AII]*
46. *In acute care, place patients who require Droplet Precautions in a single room with dedicated toilet and patient sink when available. [AII]*
47. *In long-term care and other residential settings, residents who require Droplet Precautions should remain in their room or bed space if feasible. [AII]*
48. *In ambulatory settings, offer mask and hand hygiene to client/patient at triage. Triage client/patient away from waiting area to a single room as soon as possible, or maintain a two-metre spatial separation. [AII]*
49. *Wear a mask and eye protection within two metres of a client/patient/resident on Droplet Precautions. [BII]*
50. *Whenever possible, dedicate equipment and items in the environment. [AII]*
51. *Clients/patients/residents on Droplet Precautions should wear a mask for transport or ambulation outside of their room, if tolerated. [BIII]*
52. *Clients/patients/residents who require Airborne Precautions must be moved to an airborne isolation room as soon as possible. [AII]*
53. *Clients/patients/residents on Airborne Precautions must remain in the room with the door closed, unless leaving the room for medically necessary procedures. [BII]*
54. *An N95 respirator must be worn by each person who enters an airborne isolation room when it is being used for tuberculosis. [AII]*
55. *Only immune staff may enter the room of a patient with measles, varicella or zoster. [AIII]*
56. *Clients/patients/residents who require Airborne Precautions must wear a mask during transport or activities outside their room, if tolerated. [BIII]*
57. *Transport staff should wear an N95 respirator during transport of clients/patients/residents on Airborne Precautions. [CIII]*

Occupational Health and Hygiene Issues

It has been shown that adherence of health care providers to the recommended infection prevention and control practices will decrease the transmission of infectious agents in health care settings.¹⁹ All health care providers have a responsibility to know their immunization status; to adhere to Routine Practices and Additional Precautions (including appropriate and correct use of PPE and hand hygiene); and to report exposures and infections that put themselves at risk for transmission of infections. See Section II for immunization recommendations.

POST-EXPOSURE FOLLOW-UP

The effective management of staff exposures requires the cooperation of both Occupational Health and Infection Prevention and Control staff.

Occupational health policies and procedures should address post-exposure follow-up and prophylaxis when indicated.⁶³ There should be a program to deal with staff exposures which includes¹⁰:

- a) identification of exposed staff;
- b) assessment and immunization history;
- c) post-exposure prophylaxis and follow-up including:
 - i) collection and analysis of exposures; and
 - ii) a program for prompt response to sharps injuries^{16, 63};
- d) policies to deal with spills and staff exposure to blood or body fluids; and
- e) education regarding preventive actions that may be put into place to improve practices and prevent recurrence.

RESPIRATORY PROTECTION PROGRAM, FIT-TESTING AND SEAL-CHECKING

A respiratory protection program is required for staff who will be required to wear an N95 respirator (Ministry of Labour requirement). The program must include:

- a) a health assessment;
- b) N95 respirator fit-testing; and
- c) training – health care providers and other staff required to wear an N95 respirator must be educated regarding the proper way to perform a seal-check; see [Box 10](#) for items that must be included in training.

*Fit-testing*⁴ is the use of a qualitative or quantitative method to evaluate the fit of a specific make, model and size of respirator on an individual. This procedure is to be done periodically, at least every two years and whenever there is a change in respirator face piece or the user's physical condition which could affect the respirator fit.^{4, 8, 10}

Seal-checking (also referred to as a 'fit-check') is a procedure that the health care provider must perform each time an N95 respirator is worn to ensure the respirator fits the wearer's face correctly to provide adequate respiratory protection. The health care provider is to receive training on how to perform a fit-check correctly in order to obtain a tight facial seal.^{7, 96}

Recommendations for Occupational Health and Hygiene

58. Staff who are required to wear PPE will receive instruction in the appropriate and correct use and disposal of barrier equipment. [BII]

59. The health care setting will have a program to deal with staff exposures, including exposures to blood and body fluids. [All]

60. The health care setting shall have a respiratory protection program for staff who will be required to wear an N95 respirator. [Ministry of Labour Requirement]

Audits of Compliance with Feedback

In order to achieve long-term improvement, the health care setting must make infection prevention an institutional priority and integrate infection prevention and control practices into the organization's safety culture.^{10, 18, 98} Improving adherence to infection control practices requires a multifaceted approach that incorporates ongoing education and continuous assessment of both the individual and the work environment.¹⁸ Staffing levels should be adequate to allow for compliance.⁹⁹

Compliance with Routine Practices and Additional Precautions may be related to several factors⁹⁸:

- a) perceived value of preventive actions;
- b) job hindrances (e.g., increased workload, interference with job duties, physical discomfort when wearing PPE);
- c) availability of PPE in the work area;
- d) provision of employee feedback/reinforcement with respect to adherence; and
- e) organizational level factors promoting a safety climate in the workplace.

Most strategies for the evaluation of application of Routine Practices and Additional Precautions are based on observational audits of compliance and performance feedback with recommendations for improvement. These strategies include:

- a) knowledge and application of written guidelines;
- b) correct selection and removal of PPE; and
- c) compliance with hand hygiene procedures.

Facilities where results of audits and feedback identify issues relating to compliance should provide ongoing educational and motivational activities to encourage long-lasting improvement in infection prevention and control practices.

There should be a plan of action for persistent failure. Non-compliance should not be tolerated, as this is a patient and health care provider safety issue. Compliance results should be part of the performance appraisal.

Summary of Recommendations for Routine Practices And Additional Precautions In All Health Care Settings

This summary table is intended to assist with self-assessment internal to the health care setting for quality improvement purposes.

See complete text for rationale.

Recommendation		Compliant	Partial Compliance	Non-compliant	Action Plan	Accountability
ROUTINE PRACTICES						
1.	<i>The elements of Routine Practices must be incorporated into the culture of all health care settings and into the daily practice of each health care provider during the care of all clients/patients/residents at all times. [BII]</i>					
2.	<i>Visitors should receive instruction regarding specific facility control measures before they visit a client/patient/resident, to ensure compliance with established practices. [BII]</i>					
3.	<i>Perform a risk assessment before each interaction with a client/patient/resident or their environment in order to determine which interventions are required to prevent transmission during the planned interaction. [BIII]</i>					
4.	<i>Choose client/patient/resident accommodation based on the risk assessment.</i>					
5.	<i>Choose personal protective equipment based on the risk assessment.</i>					

Recommendation		Compliant	Partial Compliance	Non-compliant	Action Plan	Accountability
6.	<i>All health care settings must implement a comprehensive hand hygiene program that follows the best practices recommended in 'Best Practices for Hand Hygiene in All Health Care Settings'.</i>					
7.	<i>Provide sufficient supplies of easily accessible PPE. [AIII]</i>					
8.	<i>Implement a process for evaluating PPE to ensure it meets quality standards where applicable, including a respiratory protection program compliant with the Ministry of Labour requirements. [AIII]</i>					
9.	<i>Provide education in the proper use of PPE to all health care providers and other staff who have the potential to be exposed to blood and body fluids. [BII]</i>					
10.	<i>Wear gloves when it is anticipated that the hands will be in contact with mucous membranes, non-intact skin, tissue, blood, body fluids, secretions, excretions, or equipment and environmental surfaces contaminated with the above. [AII]</i>					
11.	<i>Gloves are not required for routine health care activities in which contact is limited to the intact skin of the client/patient/resident. [AIII]</i>					
12.	<i>Select gloves that fit well and are of sufficient durability for the task. [AII]</i>					
13.	<i>Put on gloves just before the task or procedure that requires them. [AII]</i>					
14.	<i>Perform hand hygiene before putting on gloves for aseptic procedures.</i>					

Recommendation		Compliant	Partial Compliance	Non-compliant	Action Plan	Accountability
15.	<i>Remove gloves immediately after completion of the task that requires gloves, before touching clean environmental surfaces. [AIII]</i>					
16.	<i>Clean hands immediately after removing gloves. [AII]</i>					
17.	<i>Single-use disposable gloves should not be re-used or washed. [AII]</i>					
18.	<i>Wear a gown when it is anticipated that a procedure or care activity is likely to generate splashes or sprays of blood, body fluids, secretions, or excretions. [BIII]</i>					
19.	<i>Remove gown immediately after the task for which it has been used in a manner that prevents contamination of clothing or skin and prevents agitation of the gown. [BII]</i>					
20.	<i>Wear a mask and eye protection to protect the mucous membranes of the eyes, nose and mouth when it is anticipated that a procedure or care activity is likely to generate splashes or sprays of blood, body fluids, secretions or excretions. [AII]</i>					
21.	<i>Wear an N95 respirator to prevent inhalation of small particles that may contain infectious agents transmitted via the <u>airborne</u> route. [AII]</i>					
22.	<i>Single rooms, with dedicated bathroom and sink, are preferred for placement of all clients/patients/residents.[BII]</i>					
23.	<i>If single rooms are limited, there should be clear protocols for determining options for patient placement and room sharing based on a risk assessment. [BII]</i>					

Recommendation		Compliant	Partial Compliance	Non-compliant	Action Plan	Accountability
24.	<i>Clients/patients/residents who visibly soil the environment or for whom appropriate hygiene cannot be maintained should be placed in single rooms with dedicated toileting facilities. [AIII]</i>					
25.	<i>A sharps injury prevention program must be in place in all health care settings. [AII]</i>					
26.	<i>Appropriate policies and procedures are in place to ensure staff attendance at training/education in Routine Practices (including hand hygiene) and attendance is recorded and reported back to the manager to become a part of the employee's performance review. [AII]</i>					
27.	<i>There is a program that promotes respiratory etiquette to staff, clients/patients/residents and visitors in the health care setting. [AII]</i>					
28.	<i>There is a clear expectation that staff do not come into work when ill with symptoms that are of an infectious origin, and this expectation is supported with appropriate attendance management policies. [BII]</i>					
ADDITIONAL PRECAUTIONS						
29.	<i>The elements of Additional Precautions must be incorporated into the health care practices of each health care setting. [BII]</i>					
30.	<i>Appropriate policies and procedures are in place to ensure staff attendance at training/education in Additional Precautions and attendance is recorded and reported back to the manager to become a part of the employee's performance review. [AII]</i>					

Recommendation		Compliant	Partial Compliance	Non-compliant	Action Plan	Accountability
31.	<i>When single patient rooms are limited, determine the feasibility of cohorting patients who are infected or colonized with the same microorganism. [BIII]</i>					
32.	<i>Consider the use of geographic cohorting of patients and staff to reduce transmission during outbreaks. [AII]</i>					
33.	<i>When cohorting, Additional Precautions must be applied individually for each patient/resident within the cohort. Gowns and gloves must not be worn from patient-to-patient within the cohort and patient care equipment must not be shared.</i>					
34.	<i>Visitors to clients/patients/residents on Additional Precautions must wear the same personal protective equipment as health care providers if they will be in contact with other clients/patients/residents or are providing direct care. [BIII]</i>					
35.	<i>Each health care setting should have a policy authorizing any regulated health care professional to initiate the appropriate Additional Precautions at the onset of symptoms. [BII]</i>					
36.	<i>Additional Precautions should remain in place until there is no longer a risk of transmission of the microorganism or illness. [AII]</i>					
37.	<i>The health care setting should have a policy that permits discontinuation of Additional Precautions in consultation with the Infection Prevention and Control Professional or designate. [BIII]</i>					

Recommendation		Compliant	Partial Compliance	Non-compliant	Action Plan	Accountability
38.	<i>Additional Precautions should not be used any longer than necessary; ongoing assessment of the risk of transmission should be performed by Infection Prevention and Control Professionals.[All]</i>					
39.	<i>In acute care, place patients who require Contact Precautions in a single room with dedicated toilet and patient sink when available. [All]</i>					
40.	<i>In long-term care and other residential settings, placement of residents who require Contact Precautions should be determined on a case-by-case basis using a risk assessment. [BII]</i>					
41.	<i>In ambulatory settings, place patients who require Contact Precautions in an examination room or cubicle as soon as possible. [BII]</i>					
42.	<i>In acute care, wear gloves for all activities in the patient's room or bed space. Remove gloves and perform hand hygiene immediately on leaving the room or bed space. [All]</i>					
43.	<i>In acute care, wear a gown for all activities where skin or clothing will come in contact with the patient or the patient's environment. When indicated, put on gown on entry to the patient's room or bed space. If used, remove gown and perform hand hygiene immediately on leaving the room or bed space. [BIII]</i>					

Recommendation		Compliant	Partial Compliance	Non-compliant	Action Plan	Accountability
44.	<i>In non-acute settings, wear gloves and a gown for activities that involve direct care. Remove gloves and gown, if worn, and perform hand hygiene immediately on leaving the room. [AII]</i>					
45.	<i>Whenever possible, dedicate equipment and items to the patient/resident. [AII]</i>					
46.	<i>In acute care, place patients who require Droplet Precautions in a single room with dedicated toilet and patient sink when available. [AII]</i>					
47.	<i>In long-term care and other residential settings, residents who require Droplet Precautions should remain in their room or bed space if feasible. [AII]</i>					
48.	<i>In ambulatory settings, offer mask and hand hygiene to client/patient at triage. Triage client/patient away from waiting area to a single room as soon as possible, or maintain a two-metre spatial separation. [AII]</i>					
49.	<i>Wear a mask and eye protection within two metres of a client/patient/resident on Droplet Precautions. [BII]</i>					
50.	<i>Whenever possible, dedicate equipment and items in the environment. [AII]</i>					

Recommendation		Compliant	Partial Compliance	Non-compliant	Action Plan	Accountability
51.	<i>Clients/patients/residents on Droplet Precautions must wear a mask for transport or ambulation outside of their room, if tolerated. [BIII]</i>					
52.	<i>Clients/patients/residents who require Airborne Precautions must be moved to an airborne isolation room as soon as possible. [AII]</i>					
53.	<i>Clients/patients/residents on Airborne Precautions must remain in the room with the door closed, unless leaving the room for medically necessary procedures. [BII]</i>					
54.	<i>An N95 respirator must be worn by each person who enters an airborne isolation room when it is being used for tuberculosis. [AII]</i>					
55.	<i>Only immune staff may enter the room of a patient with measles, varicella or zoster. [AIII]</i>					
56.	<i>Clients/patients/residents who require Airborne Precautions must wear a mask during transport or activities outside their room, if tolerated. [BIII]</i>					
57.	<i>Transport staff should wear an N95 respirator during transport of clients/patients/residents on Airborne Precautions. [CIII]</i>					

Recommendation		Compliant	Partial Compliance	Non-compliant	Action Plan	Accountability
OCCUPATIONAL HEALTH AND HYGIENE ISSUES						
58.	<i>Staff who are required to wear PPE will receive instruction in the appropriate and correct use and disposal of barrier equipment. [BII]</i>					
59.	<i>The health care setting will have a program to deal with staff exposures, including exposures to blood and body fluids. [All]</i>					
60.	<i>The health care setting shall have a respiratory protection program for staff who will be required to wear an N95 respirator. [Ministry of Labour Requirement]</i>					

Appendices

Appendix A: Ranking System for Recommendations


Source: Public Health Agency of Canada

Categories for strength of each recommendation	
CATEGORY	DEFINITION
A	Good evidence to support a recommendation for use.
B	Moderate evidence to support a recommendation for use.
C	Insufficient evidence to support a recommendation for or against use
D	Moderate evidence to support a recommendation against use.
E	Good evidence to support a recommendation against use.
Categories for quality of evidence on which recommendations are made	
GRADE	DEFINITION
I	Evidence from at least one properly randomized, controlled trial.
II	Evidence from at least one well-designed clinical trial without randomization, from cohort or case-controlled analytic studies, preferably from more than one centre, from multiple time series, or from dramatic results in uncontrolled experiments.
III	Evidence from opinions of respected authorities on the basis of clinical experience, descriptive studies, or reports of expert committees.


Appendix B: Performing a Risk Assessment Related to Routine Practices and Additional Precautions

An **individual assessment** of each client/patient/resident’s potential risk of transmission of microorganisms must be made by all health care providers and other staff who come into contact with them. Based on that risk assessment and a risk assessment of the task, one may determine appropriate **intervention and interaction** strategies, such as hand hygiene, waste management, use of personal protective equipment (PPE) and client/patient/resident placement, that will reduce the risk of transmission of microorganisms to and from the individual.²⁸ When a client/patient/resident has undiagnosed symptoms or signs of an infection, interventions must be informed by **organizational requirements**.

Risk Assessment Steps to be Performed by a Health Care Provider to Determine an Individual’s Risk of Transmission of Infectious Agents and the Rationale for Associated Protective Measures



Routine Practices Risk Assessment Algorithm for All Client/Patient/Resident Interactions



Appendix C: Decision-Making Related to Accommodation and Additional Precautions

A single room is the preferred accommodation for all clients/patients/residents in all health care settings. Where single rooms are not available, the following considerations may be taken into account:

1. Accommodation for Patients/Residents Requiring Droplet Precautions

There is a requirement for spatial separation of at least two metres and facial protection for close contact with a patient/resident with a new/worse cough or shortness of breath with fever, or copious uncontrolled respiratory secretions. The following may be used to determine placement:

- Does the patient/resident have:
 - A new or worse cough or shortness of breath with fever or chills?
 - Copious uncontrolled respiratory secretions?
 - Suspected or diagnosed meningococcal disease or meningitis of unknown etiology?

If yes:



- Should be accommodated preferentially in a single room
- If a single room is not available, maintain a spatial separation of at least two metres
- Facial protection for close contact with the patient/resident
- Initiate Contact Precautions if indicated (e.g., respiratory viral infection also spread by the contact route, such as influenza)

2. Accommodation for Patients/Residents with MRSA

Patients/residents known to be colonized or infected with MRSA should be placed in a single room with individual toileting facilities. In acute care settings, MRSA-positive patients should not share rooms with MRSA-negative patients.

When single rooms for Contact Precautions are limited, priority should be given to patients/residents who are at increased risk of disseminating microorganisms into the environment:

- Does the patient/resident have:
 - A respiratory infection?
 - Colonized tracheostomy and/or uncontrolled respiratory secretions?
 - Wound or stoma drainage not contained by a dressing or appliance?
 - Desquamating skin condition (e.g., psoriasis, burns)?
 - Cognitive impairment?
 - Poor compliance with personal hygiene?

If yes:



- Should be accommodated preferentially in a single room
 - If a single room is not available, cohort with other patients/residents with MRSA, in consultation with Infection Prevention and Control and on a case-by-case basis
 - Initiate Contact Precautions
- In non-acute care MRSA residents should not share a room with:
 - Individuals who have open wounds or decubitus ulcers
 - Individuals who have urinary catheters, feeding tubes or other invasive devices
 - Individuals whose hygiene is compromised
 - Individuals who have debilitating or bed-bound conditions that require extensive 'hands-on' care
 - If patients/residents with MRSA are accommodated with patients/residents who do not have MRSA, there must be increased attention to effective environmental cleaning

3. Accommodation for Patients/Residents with VRE or *Clostridium difficile* Infection (CDI)

Patients/residents known to be colonized or infected with VRE or who have CDI should be placed in a single room with individual toileting facilities. In acute care settings, VRE-positive patients should not share rooms or toileting facilities with VRE-negative patients.

When single rooms for Contact Precautions are limited, priority should be given to patients/residents who are at increased risk of disseminating microorganisms into the environment:

- Does the patient/resident have:
 - Diarrhea not contained by diapers?
 - Faecal incontinence?
 - Wound or stoma drainage not contained by a dressing or appliance?
 - Cognitive impairment?
 - Poor compliance with personal hygiene?

If yes:



- Should be accommodated preferentially in a single room
- If a single room is not available, cohort VRE patients/residents with other patients/residents with VRE, and CDI patients/residents with other patients/residents with CDI, in consultation with Infection Prevention and Control and on a case-by-case basis
- Patients/residents with VRE or CDI should use a dedicated commode or bed pan for toileting
- Increase attention to effective environmental cleaning
- Move to a single room as soon as possible

Appendix D: Time Required for Airborne Isolation Room to Clear *M. tuberculosis*

Air Changes Per Hour and Time in Minutes Required for Removal Efficiencies of 90%, 99% or 99.9% of Airborne Contaminants









# Air Changes Per Hour	Minutes required for a removal efficiency of:		
	90%	99%	99.9%
1	138	276	414
2	69	138	207
3	46	92	138
4	35	69	104
5	28	55	83
6	23	46	69
7	20	39	59
8	17	35	52
9	15	31	46
10	14	28	41
11	13	25	38
12	12	23	35
13	11	21	32
14	10	20	30
15	9	18	28
16	9	17	26
17	8	16	24
18	8	15	23
19	7	15	22
20	7	14	21

This table is prepared according to the formula $t = (ln C_2/C_1)/(Q/V) \times 60$, which is an adaptation of the formula for the rate of purging airborne contaminants (100-Mutchler 1973) with $t_1=0$ and $C_2/C_1=1 - (\text{removal efficiency}/100)$, where:

- t₁ = initial timepoint
 - C₁ = initial concentration of contaminant
 - C₂ = final concentration of contaminants
 - Q = air flow rate (cubic feet per hour)
 - V = room volume (cubic feet)
- and: $Q + V = \text{ACH}$







Source: Members of the Ad Hoc Committee for the Guidelines for Preventing the Transmission of Tuberculosis in Canadian Health Care Facilities and Other Institutional Settings. 'Guidelines for Preventing the Transmission of Tuberculosis in Canadian Health Care Facilities and Other Institutional Settings'. *Can Commun Dis Rep.* 1996;22 Suppl 1:i-iv, 1-50, i-iv, 1-55.¹⁰⁰

Appendix E: PIDAC's Routine Practices Fact Sheet for All Health Care Settings

ROUTINE PRACTICES to be used with <u>ALL PATIENTS</u>	
	<p>Hand Hygiene</p> <p>Hand hygiene is performed using alcohol-based hand rub or soap and water:</p> <ul style="list-style-type: none"> ✓ Before and after each client/patient/resident contact ✓ Before performing invasive procedures ✓ Before preparing, handling, serving or eating food ✓ After care involving body fluids and before moving to another activity ✓ Before putting on and after taking off gloves and PPE ✓ After personal body functions (e.g., blowing one's nose) ✓ Whenever hands come into contact with secretions, excretions, blood and body fluids ✓ After contact with items in the client/patient/resident's environment
	<p>Mask and Eye Protection or Face Shield [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Protect eyes, nose and mouth during procedures and care activities likely to generate splashes or sprays of blood, body fluids, secretions or excretions. ✓ Wear within two metres of a coughing client/patient/resident.
	<p>Gown [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear a long-sleeved gown if contamination of skin or clothing is anticipated.
	<p>Gloves [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear gloves when there is a risk of hand contact with blood, body fluids, secretions, excretions, non-intact skin, mucous membranes or contaminated surfaces or objects. ✓ Wearing gloves is NOT a substitute for hand hygiene. ✓ Remove immediately after use and perform hand hygiene after removing gloves.
	<p>Environment and Equipment</p> <ul style="list-style-type: none"> ✓ All equipment that is being used by more than one client/patient/resident must be cleaned between clients/patients/residents. ✓ All high-touch surfaces in the client/patient/resident's room must be cleaned daily.
	<p>Linen and Waste</p> <ul style="list-style-type: none"> ✓ Handle soiled linen and waste carefully to prevent personal contamination and transfer to other clients/patients/residents.
	<p>Sharps Injury Prevention</p> <ul style="list-style-type: none"> ✓ NEVER RECAP USED NEEDLES. ✓ Place sharps in sharps containers. ✓ Prevent injuries from needles, scalpels and other sharp devices. ✓ Where possible, use safety-engineered medical devices.
	<p>Patient Placement/Accommodation</p> <ul style="list-style-type: none"> ✓ Use a single room for a client/patient/resident who contaminates the environment. ✓ Perform hand hygiene on leaving the room.

Images Developed By: Kevin Rostant







Appendix F: Sample Signage for Entrance to Room of a Patient Requiring Contact Precautions in Acute Care Facilities

CONTACT PRECAUTIONS – Acute Care Facilities	
	<p>Hand Hygiene as per Routine Practices</p> <p>Hand hygiene is performed:</p> <ul style="list-style-type: none"> ✓ Before and after each patient contact ✓ Before performing invasive procedures ✓ Before preparing, handling, serving or eating food ✓ After care involving body fluids and before moving to another activity ✓ Before putting on and after taking off gloves and other PPE ✓ After personal body functions (e.g., blowing one’s nose) ✓ Whenever hands come into contact with secretions, excretions, blood and body fluids ✓ After contact with items in the patient’s environment ✓ Whenever there is doubt about the necessity for doing so
	<p>Patient Placement</p> <ul style="list-style-type: none"> ✓ Single room with own toileting facilities ✓ Door may remain open ✓ Perform hand hygiene on leaving the room
	<p>Gown [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear a long-sleeved gown when entering the patient’s room or bed space if skin or clothing will come into direct contact with the patient or the patient’s environment
	<p>Gloves [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear gloves when entering the patient’s room or bed space ✓ Wearing gloves is NOT a substitute for hand hygiene. ✓ Remove gloves on leaving the room or bed space and perform hand hygiene
	<p>Environment and Equipment</p> <ul style="list-style-type: none"> ✓ Dedicate routine equipment to the patient (e.g., stethoscope, commode) ✓ Disinfect all equipment that comes out of the room ✓ All high-touch surfaces in the patient’s room must be cleaned at least daily
	<p>Visitors</p> <ul style="list-style-type: none"> ✓ Visitors must wear gloves and a long-sleeved gown if they will be in contact with other patients or will be providing <u>direct care</u>*, as required by Routine Practices ✓ Visitors must perform hand hygiene before entry and on leaving the room

* **Direct Care:** Providing hands-on care, such as bathing, washing, turning the patient, changing clothing, continence care, dressing changes, care of open wounds/lesions or toileting. Feeding and pushing a wheelchair are not classified as direct care.

Images Developed By: Kevin Rostant







Appendix G: Sample Signage for Entrance to Room of a Patient Requiring Contact Precautions in Non-acute Care Facilities

CONTACT PRECAUTIONS – Non-acute Care Facilities	
	<p>Hand Hygiene as per Routine Practices</p> <p>Hand hygiene is performed:</p> <ul style="list-style-type: none"> ✓ Before and after each resident contact ✓ Before performing invasive procedures ✓ Before preparing, handling, serving or eating food ✓ After care involving body fluids and before moving to another activity ✓ Before putting on and after taking off gloves and other PPE ✓ After personal body functions (e.g., blowing one’s nose) ✓ Whenever hands come into contact with secretions, excretions, blood and body fluids ✓ After contact with items in the resident’s environment ✓ Whenever there is doubt about the necessity for doing so ✓ Clean the resident’s hands before he/she leaves his/her room
	<p>Resident Placement</p> <ul style="list-style-type: none"> ✓ Single room with own toileting facilities if resident hygiene is poor ✓ Door may remain open ✓ Perform hand hygiene on leaving the room or bed space
	<p>Gown [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear a long-sleeved gown for <u>direct care</u>* when skin or clothing may become contaminated
	<p>Gloves [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear gloves for <u>direct care</u>* ✓ Wearing gloves is NOT a substitute for hand hygiene ✓ Remove gloves on leaving the room or bed space and perform hand hygiene
	<p>Environment and Equipment</p> <ul style="list-style-type: none"> ✓ Dedicate routine equipment to the resident if possible (e.g., stethoscope, commode) ✓ Disinfect all equipment before it is used for another resident ✓ All high-touch surfaces in the resident’s room must be cleaned at least daily
	<p>Visitors</p> <ul style="list-style-type: none"> ✓ Visitors must wear gloves and a long-sleeved gown if they will be in contact with other residents or will be providing <u>direct care</u>*, as required by Routine Practices ✓ Visitors must perform hand hygiene before entry and on leaving the room

* **Direct Care:** Providing hands-on care, such as bathing, washing, turning the resident, changing clothing, continence care, dressing changes, care of open wounds/lesions or toileting. Feeding and pushing a wheelchair are not classified as direct care.








Images Developed By: Kevin Rostant

Appendix H: Sample Signage for Entrance to Room of a Patient Requiring Droplet Precautions in All Health Care Facilities

DROPLET PRECAUTIONS – All Facilities	
	<p>Hand Hygiene as per Routine Practices</p> <p>Hand hygiene is performed:</p> <ul style="list-style-type: none"> ✓ Before and after each patient/resident contact ✓ Before performing invasive procedures ✓ Before preparing, handling, serving or eating food ✓ After care involving body fluids and before moving to another activity ✓ Before putting on and after taking off gloves and other PPE ✓ After personal body functions (e.g., blowing one’s nose) ✓ Whenever hands come into contact with secretions, excretions, blood and body fluids ✓ After contact with items in the patient/resident’s environment ✓ Whenever there is doubt about the necessity for doing so
	<p>Patient/Resident Placement</p> <ul style="list-style-type: none"> ✓ Single room with own toileting facilities if available, or maintain a spatial separation of at least 2 metres between the patient/resident and others in the room, with privacy curtain drawn ✓ Door may remain open ✓ Perform hand hygiene on leaving the room
	<p>Mask and Eye Protection or Face Shield</p> <ul style="list-style-type: none"> ✓ Wear within 2 metres of the patient/resident ✓ Remove and perform hand hygiene on leaving the room
	<p>Environment and Equipment</p> <ul style="list-style-type: none"> ✓ Dedicate routine equipment to the patient/resident (e.g., stethoscope, thermometer) ✓ Disinfect all equipment that comes out of the room ✓ All high-touch surfaces in the patient/resident’s room must be cleaned at least daily
	<p>Patient/Resident Transport</p> <ul style="list-style-type: none"> ✓ Patient/resident to wear a mask during transport
	<p>Visitors</p> <ul style="list-style-type: none"> ✓ Non-household visitors wear a mask and eye protection within 2 metres of the patient/resident ✓ Visitors must perform hand hygiene before entry and on leaving the room

Images Developed By: Kevin Rostant








Appendix I: Sample Signage for Entrance to Room of a Patient Requiring Droplet and Contact Precautions in Acute Care Facilities

DROPLET + CONTACT PRECAUTIONS – Acute Care Facilities			
	<p>Hand Hygiene as per Routine Practices</p> <p>Hand hygiene is performed:</p> <ul style="list-style-type: none"> ✓ Before and after each patient contact ✓ Before performing invasive procedures ✓ Before preparing, handling, serving or eating food ✓ After care involving body fluids and before moving to another activity ✓ Before putting on and after taking off gloves and other PPE ✓ After personal body functions (e.g., blowing one’s nose) ✓ Whenever hands come into contact with secretions, excretions, blood and body fluids ✓ After contact with items in the patient’s environment ✓ Whenever there is doubt about the necessity for doing so 		
	<p>Patient Placement</p> <ul style="list-style-type: none"> ✓ Single room with own toileting facilities if available, or maintain a spatial separation of at least 2 metres between the patient and others in the room, with privacy curtain drawn ✓ Door may remain open ✓ Perform hand hygiene on leaving the room 		
	<p>Mask and Eye Protection or Face Shield</p> <ul style="list-style-type: none"> ✓ Wear within 2 metres of the patient ✓ Remove and perform hand hygiene on leaving the room 		
	<p>Gown [based on risk assessment] and Gloves</p> <ul style="list-style-type: none"> ✓ Wear gloves when entering the patient’s room or bed space ✓ Wearing gloves is NOT a substitute for hand hygiene. ✓ Remove gloves on leaving the room or bed space and perform hand hygiene ✓ Wear a long-sleeved gown when entering the patient’s room or bed space if skin or clothing will come into direct contact with the patient or the patient’s environment 		
	<p>Environment and Equipment</p> <ul style="list-style-type: none"> ✓ Dedicate routine equipment to the patient (e.g., stethoscope, thermometer) ✓ Disinfect all equipment that comes out of the room ✓ All high-touch surfaces in the patient’s room must be cleaned at least daily 		
	<p>Patient Transport</p> <ul style="list-style-type: none"> ✓ Patient to wear a mask during transport 		<p>Visitors</p> <ul style="list-style-type: none"> ✓ Non-household visitors wear a mask and eye protection within 2 metres of the patient ✓ Visitors must wear gloves and a long-sleeved gown if they will be in contact with other patients or will be providing <u>direct care</u>* ✓ Visitors must perform hand hygiene before entry and on leaving the room

* **Direct Care:** Providing hands-on care, such as bathing, washing, turning the patient, changing clothing, continence care, dressing changes, care of open wounds/lesions or toileting. Feeding and pushing a wheelchair are not classified as direct care.







Images Developed By: Kevin Rostant

Appendix J: Sample Signage for Entrance to Room of a Resident Requiring Droplet and Contact Precautions in Non-acute Care Facilities

DROPLET + CONTACT PRECAUTIONS – Non-acute Care Facilities	
	<p>Hand Hygiene as per Routine Practices</p> <p>Hand hygiene is performed:</p> <ul style="list-style-type: none"> ✓ Before and after each resident contact ✓ Before performing invasive procedures ✓ Before preparing, handling, serving or eating food ✓ After care involving body fluids and before moving to another activity ✓ Before putting on and after taking off gloves and other PPE ✓ After personal body functions (e.g., blowing one’s nose) ✓ Whenever hands come into contact with secretions, excretions, blood and body fluids ✓ After contact with items in the resident’s environment ✓ Whenever there is doubt about the necessity for doing so
	<p>Resident Placement</p> <ul style="list-style-type: none"> ✓ Single room with own toileting facilities if resident hygiene is poor and if available, or maintain a spatial separation of at least 2 metres between the resident and others in the room, with privacy curtain drawn ✓ Door may remain open ✓ Perform hand hygiene on leaving the room
	<p>Mask and Eye Protection or Face Shield</p> <ul style="list-style-type: none"> ✓ Wear within 2 metres of the resident ✓ Remove and perform hand hygiene on leaving the room
	<p>Gown and Gloves [based on risk assessment]</p> <ul style="list-style-type: none"> ✓ Wear a long-sleeved gown for <u>direct care</u>* when skin or clothing may become contaminated ✓ Wear gloves for <u>direct care</u>* ✓ Wearing gloves is NOT a substitute for hand hygiene. ✓ Remove gloves on leaving the room or bed space and perform hand hygiene
	<p>Environment and Equipment</p> <ul style="list-style-type: none"> ✓ Dedicate routine equipment to the resident if possible (e.g., stethoscope, thermometer) ✓ Disinfect all equipment before it is used for another resident ✓ All high-touch surfaces in the patient’s room must be cleaned at least daily
	<p>Resident Transport</p> <ul style="list-style-type: none"> ✓ Resident to wear a mask during transport
	<p>Visitors</p> <ul style="list-style-type: none"> ✓ Non-household visitors wear a mask and eye protection within 2 metres of the resident ✓ Visitors must wear gloves and a long-sleeved gown if they will be in contact with other residents or will be providing <u>direct care</u>* ✓ Visitors must perform hand hygiene before entry and on leaving the room

* Direct Care: Providing hands-on care, such as bathing, washing, turning the resident, changing clothing, continence care, dressing changes, care of open wounds/lesions or toileting. Feeding and pushing a wheelchair are not classified as direct care.

Appendix K: Sample Signage for Entrance to Room of a Patient Requiring Airborne Precautions in All Health Care Facilities


AIRBORNE PRECAUTIONS – All Facilities	
	<p>Hand Hygiene as per Routine Practices</p> <p>Hand hygiene is performed:</p> <ul style="list-style-type: none"> ✓ Before and after each patient/resident contact ✓ Before performing invasive procedures ✓ Before preparing, handling, serving or eating food ✓ After care involving body fluids and before moving to another activity ✓ Before putting on and after taking off gloves and other PPE ✓ After personal body functions (e.g., blowing one’s nose) ✓ Whenever hands come into contact with secretions, excretions, blood and body fluids ✓ After contact with items in the patient/resident’s environment ✓ Whenever there is doubt about the necessity for doing so
	<p>Patient/Resident Placement</p> <ul style="list-style-type: none"> ✓ Single room with own toileting facilities ✓ Room must have negative pressure ventilation with room air exhausted outside or through a HEPA filter ✓ Monitor negative pressure daily while in use ✓ Door must remain closed
	<p>N95 Respirator</p> <ul style="list-style-type: none"> ✓ Wear a fit-tested, seal-checked N95 respirator for entry to the room for TB patients ✓ For measles, varicella or disseminated zoster, only immune staff are to enter the room and an N95 respirator is not required
	<p>Environment and Equipment</p> <ul style="list-style-type: none"> ✓ Equipment that is being used by more than one patient/resident must be cleaned between patients/residents ✓ All high-touch surfaces in the patient’s room must be cleaned at least daily
	<p>Transport of the Patient/Resident</p> <ul style="list-style-type: none"> ✓ Patient/resident to wear a mask during transport ✓ Transport staff to wear an N95 respirator during transport
	<p>Visitors</p> <ul style="list-style-type: none"> ✓ Visitors must be kept to a minimum ✓ Visitors must perform hand hygiene before entry and on leaving the room ✓ For TB, household members do not require an N95 respirator ✓ For TB, non-household visitors require an N95 respirator ✓ For measles/varicella, visitors should be counselled before entering the room

Images Developed By: Kevin Rostant

Appendix L: Recommended Steps for Putting On and Taking Off Personal Protective Equipment (PPE)

Images developed by Kevin Rostant.


Some images adapted from Northwestern Ontario Infection Control Network – NWOICN



TAKING OFF PPE

1. Remove Gloves

- Remove gloves using a glove-to-glove/skin-to-skin technique
- Grasp outside edge near the wrist and peel away, rolling the glove inside-out
- Reach under the second glove and peel away
- Discard immediately into waste receptacle



2. Remove Gown

- Remove gown in a manner that prevents contamination of clothing or skin
- Starting at the neck ties, the outer, 'contaminated', side of the gown is pulled forward and turned inward, rolled off the arms into a bundle, then discarded immediately in a manner that minimizes air disturbance



6. Perform Hand Hygiene




3. Perform Hand Hygiene



5. Remove Mask/N95 Respirator

- Ties/ear loops/straps are considered 'clean' and may be touched with hands
- The front of the mask/respirator is considered to be contaminated
- Untie bottom tie then top tie, or grasp straps or ear loops
- Pull forward off the head, bending forward to allow mask/respirator to fall away from the face
- Discard immediately into waste receptacle



4. Remove Eye Protection

- Arms of goggles and headband of face shields are considered to be 'clean' and may be touched with the hands
- The front of goggles/face shield is considered to be contaminated
- Remove eye protection by handling ear loops, sides or back only
- Discard into waste receptacle or into appropriate container to be sent for reprocessing
- Personally-owned eyewear may be cleaned by the individual after each use



Appendix M: Advantages and Disadvantages of Barrier Equipment

MEDICAL GLOVES

Type	Use	Advantages	Disadvantages
Vinyl	<ul style="list-style-type: none"> ▪ Protection for: <ul style="list-style-type: none"> ○ Minimal exposure to blood/body fluids/infectious agents ○ Contact with strong acids and bases, salts, alcohols ○ Short duration tasks ▪ Protection for staff with documented skin breakdown 	<ul style="list-style-type: none"> ▪ Good level of protection but based on the quality of manufacturer ▪ Medium chemical resistance 	<ul style="list-style-type: none"> ▪ Not recommended for contact with solvents, aldehydes, ketones ▪ Quality varies with manufacturers ▪ Punctures easily when stressed ▪ Rigid – non elastic
Latex	<ul style="list-style-type: none"> ▪ Activities that require sterility ▪ Protection for: <ul style="list-style-type: none"> ○ Heavy exposure to blood/body fluids/infectious agents ○ Contact with weak acids and bases, alcohols 	<ul style="list-style-type: none"> ▪ Good barrier qualities ▪ Strong and durable ▪ Has re-seal qualities ▪ Good comfort and fit ▪ Good protection from most caustics and detergents 	<ul style="list-style-type: none"> ▪ Not recommended for contact with oils, greases and organics ▪ Contraindicated for individuals who have allergic reactions or sensitivity to latex
Nitrile	<ul style="list-style-type: none"> ▪ Protection for: <ul style="list-style-type: none"> ○ Heavy exposure to blood/body fluids/infectious agents ○ Tasks of longer duration ○ Tasks with high stress on glove ○ Tasks requiring additional dexterity ○ Chemicals and chemotherapeutic agents ○ Recommended for contact with oils, greases, acids, bases ○ Sensitivity to vinyl ▪ Preferred replacement for vinyl gloves when a documented allergy or sensitivity occurs 	<ul style="list-style-type: none"> ▪ Offers good dexterity ▪ Strong and durable ▪ Puncture-resistant ▪ Good comfort and fit ▪ Excellent resistance to chemicals 	<ul style="list-style-type: none"> ▪ Not recommended for contact with solvents, ketones, esters
Neoprene	<ul style="list-style-type: none"> ▪ Replacement sterile glove for latex when a documented allergy or sensitivity occurs ▪ Recommended for contact with acids, bases, alcohols, fats, oils, phenol, glycol ethers 	<ul style="list-style-type: none"> ▪ Good barrier qualities ▪ Strong and durable ▪ Good comfort and fit ▪ Good protection from caustics 	<ul style="list-style-type: none"> ▪ Not recommended for contact with solvents

Adapted from Sunnybrook Health Sciences Centre, Patient Care Policy Manual Section II: Infection Prevention and Control [Policy No: II-D-1200, 'Gloves'. Revised July, 2007 and London Health Sciences Centre, Occupational Health and Safety Services, 'Glove Selection and Use'. Revised April 26, 2005.

MASKS AND N95 RESPIRATORS

Type of Mask	Use	Advantages	Disadvantages
Standard Face Mask ('procedure' mask or 'isolation' mask)	<ul style="list-style-type: none"> ▪ Protection for: <ul style="list-style-type: none"> ○ Minimal exposure to infectious droplets ○ Short duration tasks ○ Tasks that do not involve exposure to blood/body fluids ▪ Protection from client/patient/resident during transportation outside of room 	<ul style="list-style-type: none"> ▪ Inexpensive 	<ul style="list-style-type: none"> ▪ Not fluid or water resistant
Fluid Resistant Mask	<ul style="list-style-type: none"> ▪ Protection for: <ul style="list-style-type: none"> ○ Heavy exposure to infectious droplets or blood/body fluids 	<ul style="list-style-type: none"> ▪ Good comfort and fit ▪ Fluid resistant 	<ul style="list-style-type: none"> ▪ Expensive
Surgical Mask	<ul style="list-style-type: none"> ▪ Protection for: <ul style="list-style-type: none"> ○ Exposure to infectious droplets or blood/body fluids ○ Long duration tasks 	<ul style="list-style-type: none"> ▪ Good comfort and fit ▪ Fluid resistant ▪ Inexpensive 	
NIOSH- certified N95 respirator	<ul style="list-style-type: none"> ▪ Protection for airborne pathogens 	<ul style="list-style-type: none"> ▪ Provides protection from small particle aerosols ▪ Better face seal prevents leakage around mask 	<ul style="list-style-type: none"> ▪ Requires fit-testing, training and seal-checking ▪ Expensive ▪ Uncomfortable for long periods of use

EYE PROTECTION

Type of Eyewear	Use	Advantages	Disadvantages
Safety Glasses	<ul style="list-style-type: none"> ▪ Protection for: <ul style="list-style-type: none"> ○ Exposure to infectious droplets or blood/body fluids 	<ul style="list-style-type: none"> ▪ may be cleaned and re-used until visibility is compromised ▪ may be worn over prescription eyeglasses ▪ good visibility 	<ul style="list-style-type: none"> ▪ with continued use, visibility may be compromised
Goggles	<ul style="list-style-type: none"> ▪ Protection for: <ul style="list-style-type: none"> ○ Exposure to infectious droplets or blood/body fluids 	<ul style="list-style-type: none"> ▪ may be cleaned and re-used until visibility is compromised ▪ may be worn over prescription eyeglasses 	<ul style="list-style-type: none"> ▪ poor visibility
Face Shield	<ul style="list-style-type: none"> ▪ Protection for: <ul style="list-style-type: none"> ○ Exposure to infectious droplets or blood/body fluids 	<ul style="list-style-type: none"> ▪ may be worn over prescription eyeglasses ▪ good visibility 	
Visor attached to Mask	<ul style="list-style-type: none"> ▪ Protection for: <ul style="list-style-type: none"> ○ Minimal exposure to infectious droplets or blood/body fluids 	<ul style="list-style-type: none"> ▪ May be worn with prescription eyeglasses ▪ Quick to put on 	

Appendix N: Clinical Syndromes and Conditions with Required Level of Precautions

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
ABSCCESS	Minor	RP	No		If community-associated MRSA is suspected, use Contact Precautions until ruled out.
	Major (drainage not contained by dressing)	Contact	Yes	Continue precautions for duration of uncontained drainage.	
ADENOVIRUS INFECTION	Conjunctivitis	Contact	Yes	Continue precautions for duration of symptoms.	May cohort patients in outbreaks.
	Pneumonia	Droplet + Contact	Yes		
AIDS	See HIV				
AMOEBIASIS (Dysentery) <i>Entamoeba histolytica</i>	Adult	RP	No		Reportable Disease
	Paediatric* and incontinent or non-compliant adult	Contact	Yes		
ANTHRAX <i>Bacillus anthracis</i>	Cutaneous or pulmonary	RP	No		Reportable Disease Notify Infection Control
ANTIBIOTIC-RESISTANT ORGANISMS (AROs) - not listed elsewhere		Contact may be indicated	May be indicated	Precautions, if required, are initiated and discontinued by Infection Control.	See also listings under MRSA, VRE, ESBL and CRE.
ARTHROPOD-BORNE VIRAL INFECTIONS Eastern, Western, & Venezuelan equine encephalomyelitis; St. Louis & California encephalitis; West Nile virus		RP	No		Reportable Disease No person-to-person transmission.
ASCARIASIS (Roundworm) <i>Ascaris lumbricoides</i>		RP	No		No person-to-person transmission.
ASPERGILLOSIS <i>Aspergillus</i> species		RP	No		If several cases occur in close proximity, look for environmental source.
BABESIOSIS		RP	No		Tick-borne. Not transmitted from person-to-person except by transfusion.
BLASTOMYCOSIS <i>Blastomyces dermatitidis</i>	Cutaneous or pulmonary	RP	No		No person-to-person transmission.
BOTULISM	See Food Poisoning/Food-borne Illness				

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
BRONCHITIS/ BRONCHIOLITIS	See Respiratory Infections				
BRUCELLOSIS (Undulant fever)		RP	No		Reportable Disease No person-to-person transmission If lesions present, see Abscess
CAMPYLOBACTER	Adult	RP	No		Reportable Disease Notify Infection Control
	Paediatric* and incontinent or non-compliant adult	Contact	Yes	Continue precautions until stools are formed.	
CARBAPENEM-RESISTANT ENTEROBACTERIACEAE (CRE)	See Enterobacteriaceae, Resistant				
CAT-SCRATCH FEVER <i>Bartonella henselae</i>		RP	No		No person-to-person transmission.
CELLULITIS , with drainage	See Abscess				
CELLULITIS	Child < 5 years of age if <i>Haemophilus influenzae</i> type B is present or suspected	Droplet	Yes	Continue precautions until 24 hours of appropriate antimicrobial therapy or until <i>H. influenzae</i> type B is ruled out.	
CHANCROID <i>Haemophilus ducreyi</i>		RP	No		Reportable Disease
CHICKENPOX	See Varicella				
CHLAMYDIA	<i>Chlamydia trachomatis</i> genital infection or lymphogranuloma venereum	RP	No		Reportable Disease
	<i>Chlamydia pneumoniae, psittaci</i>	RP	No		
CHOLERA <i>Vibrio cholera</i>	Adult	RP	No		Reportable Disease Notify Infection Control
	Paediatric* and incontinent or non-compliant adult	Contact	Yes		
CLOSTRIDIUM DIFFICILE		Contact	Yes	Continue precautions until formed stool for at least two consecutive days.	Outbreaks Reportable Notify Infection Control. Laboratory-confirmed cases may be cohorted.
COCCIDIOIDOMYCOSIS (Valley Fever)	Draining lesions or pneumonia	RP	No		No person-to-person transmission.
COMMON COLD Rhinovirus		Droplet + Contact	Yes	Continue precautions for duration of symptoms.	

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
CONGENITAL RUBELLA	See Rubella				
CONJUNCTIVITIS		Contact	Yes	Continue precautions until viral aetiology ruled out or for duration of symptoms.	
COXSACKIEVIRUS	See Enteroviral Infections				
CREUTZFELDT-JAKOB DISEASE (CJD)		RP	No		Reportable Disease. Notify Infection Control. Equipment in contact with infectious material requires special handling & disinfection practices.
CROUP		Droplet + Contact	Yes	Continue precautions for duration of illness or until infectious cause ruled out.	
CRYPTOCOCCOSIS <i>Cryptococcus neoformans</i>		RP	No		No person-to-person transmission.
CRYPTOSPORIDIOSIS	Adult	RP	No		Reportable Disease Notify Infection Control
	Paediatric* and incontinent or non-compliant adult	Contact	Yes		
CYSTICERCOSIS		RP	No		No person-to-person transmission.
CYTOMEGALOVIRUS (CMV)		RP	No		Reportable Disease if congenital Transmitted by close, direct personal contact, blood transfusions or transplants.
DECUBITUS ULCER, infected	See Abscess				
DENGUE	See Arthropod-borne viral infections				
DERMATITIS		RP	Yes, if extensive		If compatible with scabies, see <i>Scabies</i>
DIARRHEA	Acute infectious	See Gastroenteritis			
	Suspected <i>C. difficile</i> diarrhea	See <i>Clostridium difficile</i>			
DIPHTHERIA <i>Corynebacterium diphtheriae</i>	Pharyngeal	Droplet	Yes	Continue precautions until two appropriate cultures taken at least 24 hours apart after cessation of antibiotics are negative for <i>C. diphtheriae</i> .	Reportable Disease Notify Infection Control
	Cutaneous	Contact	Yes		
EBOLA VIRUS	See Haemorrhagic Fevers				

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
ECHINOCOCCOSIS		RP	No		No person-to-person transmission.
ECHOVIRUS DISEASE	See Enteroviral Infections				
ENCEPHALITIS	Adult	RP	No		Reportable Disease
	Paediatric*	Contact	Yes	Continue precautions until Enterovirus is ruled out.	
ENTEROBACTERIACEAE-RESISTANT Carbapenem-resistant <i>Enterobacteriaceae</i> (CRE)		Contact	Yes	Continue precautions for duration of hospitalization	Notify Infection Control If readmitted, use Contact precautions
Extended-spectrum Beta-lactamase producing <i>Enterobacteriaceae</i> (ESBL)		Contact may be indicated	May be indicated	Precautions, if indicated, are initiated and discontinued by Infection Control	Notify Infection Control
ENTEROBIASIS (Pinworm disease) <i>Enterobius vermicularis</i>		RP	No		Transmission is faecal-oral directly or indirectly through contaminated articles e.g., bedding.
ENTEROCOLITIS	See Gastroenteritis - Necrotizing Enterocolitis				
ENTEROVIRAL INFECTIONS (Coxsackie viruses, Echo viruses)	Adult	RP	No		
	Paediatric*	Contact	Yes	Continue precautions for duration of illness.	
EPIGLOTTITIS, due to <i>Haemophilus influenzae</i> Type B	Adult	RP	No		Type B is Reportable Disease.
	Paediatric*	Droplet	Yes	Continue precautions for 24 hours after start of effective therapy.	Notify Infection Control
EHRlichiosis <i>Ehrlichia chaffeensis</i>		RP	No		Tick-borne
EPSTEIN-BARR VIRUS (Infectious Mononucleosis)		RP	No		Transmitted via intimate contact with oral secretions or articles contaminated by them.
ERYSIPELAS	See Streptococcal Disease				
ERYTHEMA INFECTIOSUM (Parvovirus B19)	Aplastic crisis	Droplet	Yes	Continue precautions for duration of hospitalization with immunocompromised persons, or 7 days with others.	
	Fifth disease	RP	No		No longer infectious by the time rash appears.

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
ESCHERICHIA COLI O157:H7	Adult	RP	No		Reportable Disease Notify Infection Control
	Paediatric* and incontinent or non-compliant adult	Contact	Yes	Continue precautions until stools are formed.	
EXTENDED SPECTRUM BETA-LACTAMASE-PRODUCING ENTEROBACTERIACEAE (ESBL)	See <i>Enterobacteriaceae</i> , Resistant				
FIFTH DISEASE	See <i>Erythema Infectiosum</i>				
FOOD POISONING / FOOD-BORNE ILLNESS	<i>Clostridium botulinum</i> (Botulism)	RP	No		Reportable Disease No person-to-person transmission.
	<i>Clostridium perfringens</i>	RP	No		
	Salmonella or <i>Escherichia coli</i> O157:H7 in paediatric or incontinent adult if stool cannot be contained	Contact	Yes	Continue precautions until Salmonellosis or <i>E. coli</i> O157:H7 are ruled out.	Reportable Disease Notify Infection Control
	Other causes	RP	No		
FRANCISELLA TULARENSIS	See Tularemia				
FURUNCULOSIS <i>Staphylococcus aureus</i>	See Abscess				
GANGRENE	Gas gangrene due to any bacteria	RP	No		No person-to-person transmission.
GASTROENTERITIS	Acute infectious	Contact	Yes	Continue precautions until <i>C.difficile</i> and Norovirus or other viral agents ruled out.	Outbreaks are reportable Notify Infection Control See specific organism if identified.
	Paediatric* and incontinent/non-compliant adult	Contact	Yes	Continue precautions for duration of illness.	
GERMAN MEASLES	See Rubella				
GIARDIASIS <i>Giardia lamblia</i>	Adult	RP	No		Reportable Disease
	Paediatric* and incontinent or non-compliant adult	Contact	Yes	Continue precautions until stools are formed	
GONORRHEA <i>Neisseria gonorrhoeae</i>		RP	No		Reportable Disease Sexual transmission.
GRANULOMA INGUINALE		RP	No		Sexual transmission.

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
HAEMOPHILUS INFLUENZAE TYPE B	Pneumonia - adult	RP	No		Reportable Disease if invasive
	Pneumonia – paediatric*	Droplet	Yes	Continue precautions until 24 hours after effective treatment	
	Meningitis	See Meningitis			
HAND, FOOT, & MOUTH DISEASE	See Enteroviral Infection				
HANTAVIRUS PULMONARY SYNDROME		RP	No		Reportable Disease No person-to-person transmission.
HANSEN'S DISEASE	See Leprosy				
HAEMORRHAGIC FEVERS (e.g., Lassa, Ebola, Marburg)		Droplet + Contact Airborne if pneumonia	Yes, with negative air flow, door closed if pneumonia	Continue precautions until symptoms resolve	Notify Public Health immediately Notify Infection Control immediately
HEPATITIS, VIRAL Hepatitis A & E	Adult	RP	No		Reportable Disease
	Paediatric* and incontinent or non- compliant adult	Contact	Yes	Duration of precautions: < 3years: duration of hospital stay > 3years: one week from symptoms onset	
Hepatitis B & C (including Delta)		RP	No		Reportable Disease Report to Occupational Health if health care provider has percutaneous or mucous membrane exposure
HERPANGINA	See Enterovirus				
HERPES SIMPLEX	Encephalitis	RP	No		Reportable Disease
	Mucocutaneous - recurrent	RP	No		
	Disseminated/ severe	Contact	Yes	Continue precautions until lesions crusted and dry.	Reportable Disease Notify Infection Control
	Neonatal infection, and infants born to mothers with active genital herpes until neonatal infection ruled out	Contact		Continue precautions for duration of symptoms	
HISTOPLASMOSIS <i>Histoplasma capsulatum</i>		RP	No		No person-to-person transmission.

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
HIV		RP	No		Reportable Disease Report to Occupational Health if health care provider has percutaneous or mucous membrane exposure
HOOKWORM DISEASE (Ancylostomiasis)		RP	No		No person-to-person transmission.
HUMAN HERPESVIRUS 6 (Roseola)	See Roseola				
IMPETIGO	See Abscess				
INFECTIOUS MONONUCLEOSIS	See Epstein-Barr virus				
INFLUENZA (seasonal)		Droplet + Contact	Yes	Continue precautions for 5 days after onset of illness.	Reportable Disease Notify Infection Control
KAWASAKI SYNDROME		RP	No		
LASSA FEVER	See Haemorrhagic Fevers				
LEGIONNAIRES' DISEASE <i>Legionella pneumophila</i>		RP	No		Reportable Disease Notify Infection Control No person-to-person transmission.
LEPROSY (Hansen's disease) <i>Mycobacterium leprae</i>		RP	No		Reportable Disease
LEPTOSPIROSIS <i>Leptospira</i> sp.		RP	No		No person-to-person transmission.
LICE	See Pediculosis				
LISTERIOSIS <i>Listeria monocytogenes</i>		RP	No		Reportable Disease
LYME DISEASE <i>Borrelia burgdorferi</i>		RP	No		Reportable Disease No person-to-person transmission.
LYMPHOCYTIC CHORIOMENINGITIS (Aseptic meningitis)		RP	No		No person-to-person transmission.
LYMPHOGRANULOMA VENEREUM	See Chlamydia trachomatis				

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
MALARIA <i>Plasmodium</i> species		RP	No		Reportable Disease No person-to-person transmission, except by blood transfusion.
MARBURG VIRUS	See Haemorrhagic Fevers				
MEASLES (Rubeola)		Airborne	Yes, with negative air flow, door closed	Continue precautions for four days after start of rash, and for duration of illness in immunocompromised patients.	Reportable Disease Notify Infection Control. Only immune staff should enter the room.
MENINGITIS	Aetiology unknown - adult	Droplet	Yes		Reportable Disease
	Aetiology unknown – paediatric*	Droplet + Contact	Yes		
	<i>Haemophilus influenzae</i> type B - adult	RP	No		
	<i>Haemophilus influenzae</i> type B – paediatric*	Droplet	Yes	Continue precautions for 24 hours after start of effective therapy.	
	Meningococcal (<i>Neisseria meningitidis</i>)	Droplet	Yes	Continue precautions for 24 hours after start of effective therapy.	Reportable Disease Notify Infection Control
	Other bacterial	RP	No		Reportable Disease See listings by bacterial type.
	Viral - adult (“aseptic”)	RP	No		Reportable Disease See also Enteroviral
	Viral - paediatric*	Contact	Yes		
MENINGOCOCCAL DISEASE <i>Neisseria meningitidis</i>		Droplet	Yes	Continue precautions for 24 hours after start of effective therapy.	Reportable Disease Notify Infection Control
MRSA Methicillin-resistant <i>Staphylococcus aureus</i>		Contact (+ Droplet if in sputum and coughing)	Yes	Continue precautions until discontinued by Infection Control.	
MUMPS (Infectious parotitis)		Droplet	Yes	Continue precautions for five days after onset of swelling.	Reportable Disease Notify Infection Control
MYCOBACTERIA Nontuberculosis, atypical eg., <i>Mycobacterium avium</i>		RP	No		No person-to-person transmission.
MYCOBACTERIA TUBERCULOSIS	See Tuberculosis				

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
MYCOPLASMA PNEUMONIA		Droplet	Yes	Continue precautions for duration of illness.	
NECROTIZING ENTEROCOLITIS		RP	No		Cohorting ill infants + Contact Precautions may be indicated for clusters/outbreaks. Unknown if transmissible.
NEISSERIA MENINGITIDIS	See Meningococcal Disease				
NOROVIRUS		Contact	Yes	Continue precautions until 48 hours after resolution of symptoms.	Outbreaks Reportable Notify Infection Control
OPHTHALMIA NEONATORUM	See Conjunctivitis				
PARAINFLUENZA VIRUS		Droplet + Contact	Yes	Continue precautions for duration of symptoms.	Cohorting may be necessary during outbreaks.
PARATYPHOID FEVER <i>Salmonella paratyphi</i>		RP	No		Reportable Disease
PARVOVIRUS B19	See Erythema Infectiosum				
PEDICULOSIS (Lice)		RP, plus gloves for direct patient contact	No	Continue precautions for 24 hours after application of pediculicide.	
PERTUSSIS (Whooping Cough) <i>Bordetella pertussis</i>		Droplet	Yes	Continue precautions for five days after start of treatment or three weeks if not treated.	Reportable Disease Notify Infection Control
PINWORMS	See Enterobiasis				
PLAGUE <i>Yersinia pestis</i>	Pneumonic	Droplet	Yes	Continue precautions for 48 hours of effective therapy.	Reportable Disease Notify Infection Control
	Bubonic	RP	No		
PLEURODYNIA	See Enteroviral Infection				
PNEUMONIA Aetiology unknown		Droplet + Contact	Yes	Continue precautions until aetiology established or clinical improvement on empiric therapy	
POLIOMYELITIS		Contact	Yes	Continue precautions for 6 weeks after onset of illness	Reportable Disease Notify Infection Control
PSEUDOMEMBRANOUS COLITIS	See <i>Clostridium difficile</i>				

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
PSITTACOSIS (Ornithosis) <i>Chlamydia psittaci</i>	See Chlamydia				
PHARYNGITIS	Adult	RP	No		
	Paediatric*	Droplet + Contact	Yes	Continue precautions for duration of illness, or 24 hours of effective therapy if Group A streptococcus	
Q FEVER <i>Coxiella burnetii</i>		RP	No		Reportable Disease No person-to-person transmission
RABIES Rhabdovirus		RP	No		Reportable Disease Notify Infection Control Person-to-person transmission not documented except via corneal transplantation. Open wound/mucous membrane exposure to saliva of a patient should be considered for prophylaxis
RESISTANT ORGANISMS	See Antibiotic-Resistant Organisms				
RESPIRATORY INFECTIONS , acute febrile		Droplet + Contact	Yes	Continue precautions until symptoms improve or infectious cause identified.	See specific organism, if identified.
RESPIRATORY SYNCYTIAL VIRUS (RSV)		Droplet + Contact	Yes	Continue precautions for duration of illness.	
REYE'S SYNDROME		RP	No		May be associated with viral infection.
RHEUMATIC FEVER		RP	No		Complication of a Group A streptococcal infection.
RHINOVIRUS	See Common Cold				
RINGWORM	See Tinea				
ROSEOLA INFANTUM (Exanthem Subitum, Sixth disease, HHV6)		RP	No		Transmission requires close, direct personal contact.
ROTA VIRUS		Contact	Yes	Continue precautions until formed stool.	
ROUNDWORM	See Ascariasis				

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
RUBELLA (German Measles)	Acquired	Droplet	Yes	Continue precautions for seven days after onset of rash.	Reportable Disease Notify Infection Control Only immune staff should provide care. Pregnant health care providers should <u>not</u> provide care regardless of immune status.
	Congenital	Droplet + Contact	Yes	Continue precautions for one year after birth, unless urine and nasopharyngeal cultures done after three months of age are negative.	
SALMONELLOSIS <i>Salmonella</i> species	Adult	RP	No		Reportable Disease Notify Infection Control
	Paediatric* and incontinent or non-compliant adult	Contact	Yes	Continue precautions until formed stool.	
SEVERE ACUTE RESPIRATORY SYNDROME (SARS) or Acute Respiratory Illness with travel to a high risk geographical area		Droplet + Contact N95 respirator for aerosol-generating procedures	Yes	Continue precautions 10 days following resolution of fever if respiratory symptoms have also resolved.	Reportable Disease Notify Public Health <u>immediately</u> Notify Infection Control <u>immediately</u>
SCABIES <i>Sarcoptes scabiei</i>	Limited, 'typical'	RP, gloves for skin contact	No	Continue precautions until 24 hours after application of scabicide.	
	Crusted, 'Norwegian'	Contact	Yes		
SCALDED SKIN SYNDROME	See Abscess, major				
SHIGELLOSIS <i>Shigella</i> species	See Gastroenteritis				
SHINGLES	See Varicella Zoster				
SMALLPOX	See Variola				
STAPHYLOCOCCAL DISEASE <i>Staphylococcus aureus</i>	Food poisoning	See Food Poisoning/Food-borne Illness			
	Skin, wound, or burn infection	See Abscess			
	Pneumonia - adult	RP	No		
	Pneumonia – paediatric*	Droplet	Yes	Continue precautions until 24 hours of effective therapy.	
	Toxic shock syndrome (TSS)	RP	No		
STREPTOCOCCAL DISEASE Group A <i>Streptococcus</i>	Skin, wound or burn infection, including necrotizing fasciitis	Droplet + Contact	Yes	Continue precautions until 24 hours of effective treatment.	Reportable Disease if invasive Notify Infection Control
	Toxic shock-like syndrome (TSLs)	Droplet + Contact	Yes		
	Pneumonia	Droplet	Yes		

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
STREPTOCOCCAL DISEASE, con't.	Pharyngitis/scarlet fever – paediatric*	Droplet	Yes		
	Endometritis (Puerperal Sepsis)	RP	No		
	Pharyngitis/ scarlet fever - adult	RP	No		
Group B Streptococcus	Neonatal	RP	No		Reportable Disease Notify Infection Control
<i>Streptococcus pneumonia</i> ('pneumococcus')		RP	No		
STRONGYLOIDIASIS <i>Strongyloides stercoralis</i>		RP	No		May cause disseminated disease in immunocompromised.
SYPHILIS <i>Treponema pallidum</i>		RP, gloves for contact with skin lesions	No		Reportable Disease
TAPEWORM DISEASE <i>Diphyllobothrium latum</i> (fish) <i>Hymenolepis nana</i> , <i>Taenia saginata</i> (beef) <i>Taenia solium</i> (pork)		RP	No		Autoinfection possible.
TETANUS <i>Clostridium tetani</i>		RP	No		Reportable disease No person-to-person transmission.
TINEA (Fungus infection dermatophytosis, dermatomycosis, ringworm)		RP	No		Thorough cleaning of bath and shower after use. No shared combs or brushes.
TOXOPLASMOSIS <i>Toxoplasma gondii</i>		RP	No		No person-to-person transmission except vertical.
TOXIC SHOCK SYNDROME	See Staphylococcal & Streptococcal Disease				
TRENCHMOUTH	See Vincent's angina				
TRICHINOSIS <i>Trichinella spiralis</i>		RP	No		Reportable Disease No person-to-person transmission.
TRICHOMONIASIS <i>Trichomonas vaginalis</i>		RP	No		Sexual transmission
TUBERCULOSIS <i>Mycobacterium tuberculosis</i>	Extrapulmonary, no draining lesions	RP	No		Reportable Disease Notify Infection Control Assess for concurrent pulmonary TB.

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
TUBERCULOSIS, con't.	Extrapulmonary, draining lesions	Airborne	Yes, with negative air flow and door closed	Continue precautions until drainage ceased or three consecutive negative AFB smears.	
	Pulmonary - confirmed or suspected or laryngeal disease	Airborne	Yes, with negative air flow and door closed	Continue precautions until TB ruled out. If confirmed, until patient has received two weeks of effective therapy, is improving clinically and has three consecutive sputum smears negative for AFB, collected 24 hours apart. If multidrug-resistant TB, until culture negative.	Reportable Disease Notify Infection Control
	Skin-test positive with no evidence of current disease	RP	No		Latent tuberculous infection (LTBI).
TULAREMIA <i>Francisella tularensis</i>		RP	No		Reportable Disease No person-to-person transmission. Notify Microbiology laboratory if suspected, as aerosols from cultures are infectious.
TYPHOID FEVER <i>Salmonella typhi</i>		RP	No		Reportable Disease
TYPHUS <i>Rickettsia species</i>		RP	No		Transmitted through close personal contact, but not in absence of lice.
URINARY TRACT INFECTION		RP	No		
VANCOMYCIN-RESISTANT ENTEROCOCCUS (VRE)	See VRE				
VANCOMYCIN-RESISTANT STAPHYLOCOCCUS AUREUS (VRSA)	See VRSA				
VARICELLA (Chickenpox)		Airborne	Yes, with negative air flow and door closed	Continue precautions until all vesicles have crusted and for at least five days.	Reportable Disease Notify Infection Control Neonates born to mothers with active varicella should be isolated at birth. Only immune staff should enter the room.

ORGANISM/ DISEASE	CATEGORY *	TYPE OF PRECAUTION	SINGLE ROOM?	DURATION OF PRECAUTIONS	COMMENTS
<p>* = Paediatric precautions apply to children who are incontinent or too immature to comply with hygiene</p> <p>RP = Routine Practices</p>					
VARICELLA ZOSTER (Shingles, Zoster) <i>Herpes zoster</i>	Immunocompromised patient, or disseminated	Airborne	Yes, with negative air flow and door closed	Continue precautions until all lesions have crusted and dried.	Notify Infection Control. Only immune staff should enter the room.
	Localized in all other patients	RP	No		Roommates and staff must be immune to chickenpox.
VARIOLA (Smallpox)		Airborne + Contact	Yes, with negative air flow and door closed	Continue precautions until all lesions have crusted and separated (3 to 4 weeks)	Report to Public Health <u>immediately</u> Notify Infection Control <u>immediately</u>
VIBRIO	See Gastroenteritis or Cholera				
VINCENT'S ANGINA (Trench mouth)		RP	No		
VIRAL DISEASES - Respiratory (if not covered elsewhere)		Droplet + Contact	Yes		See also specific disease/organism.
VRE Vancomycin-resistant enterococcus		Contact	Yes	Continue precautions until discontinued by Infection Control.	Notify Infection Control
VRSA Vancomycin-resistant <i>Staphylococcus aureus</i>		Contact	Yes	Continue precautions for duration of hospital stay.	Notify Infection Control
WEST NILE VIRUS (WNV)	See Arthropod-borne Viral Fevers				
WHOOPING COUGH	See Pertussis				
WOUND INFECTIONS	See Abscess				
YELLOW FEVER	See Arthropod-borne Viral Fevers				
YERSINIA ENTEROCOLITICA	See Gastroenteritis				
YERSINIA PESTIS	See Plague				
ZOSTER	See Herpes Zoster				

Based on Health Canada's 'Routine Practices and Additional Precautions for Preventing the Transmission of Infection in Health Care'⁷ and the Center for Disease Control's '2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Health Care Settings'¹⁸

References

1. Sung JJ, Yu I, Zhong NS, Tsoi K. Super-spreading events of SARS in a hospital setting: who, when, and why? *Hong Kong Med J*. 2009 Dec;15 Suppl 8:29-33.
2. Fowler RA, Guest CB, Lapinsky SE, Sibbald WJ, Louie M, Tang P, et al. Transmission of severe acute respiratory syndrome during intubation and mechanical ventilation. *Am J Respir Crit Care Med*. 2004 Jun 1;169(11):1198-202.
3. Conly JM. Personal protective equipment for preventing respiratory infections: what have we really learned? *CMAJ*. 2006 Aug 1;175(3):263.
4. Canadian Standards Association. CAN/CSA-Z94.4-02 (R2007) Selection, Use, and Care of Respirators: Occupational Health & Safety. Rexdale, Ont.: Canadian Standards Association; 2002 [cited April 6, 2011]; 103]. Available from: <http://ohsviewaccess.csa.ca/viewStandards.asp>.
5. Provincial Infectious Diseases Advisory Committee (PIDAC). Best Practices for Surveillance of Health Care-Associated Infections in Patient and Resident Populations 2008 [cited November 24, 2008]; Available from: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/surveillance-of-health-care-associated-infections.html>.
6. Canadian Standards Association. CAN/CSA Z317.13-07 Infection Control during Construction, Renovation and Maintenance of Health Care Facilities. Mississauga, Ont.: Canadian Standards Association; 2007.
7. Health Canada. Infection Control Guidelines: Routine practices and additional precautions for preventing the transmission of infection in health care [under revision]. *Can Commun Dis Rep*. 1999 Jul;25 Suppl 4:1-142.
8. Jensen PA, Lambert LA, Iademarco MF, Ridzon R. Guidelines for preventing the transmission of Mycobacterium tuberculosis in health-care settings, 2005. *MMWR Recomm Rep*. 2005 Dec 30;54(17):1-141.
9. Provincial Infectious Diseases Advisory Committee (PIDAC). Best Practices for Environmental Cleaning for Prevention and Control of Infections in All Health Care Settings. 2009 [cited June 14, 2011]; Available from: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/environmental-cleaning-for-prevention-and-control-of-infections.html>.
10. Provincial Infectious Diseases Advisory Committee (PIDAC). Best Practices for Infection Prevention and Control Programs in Ontario In All Health Care Settings 2011 [cited June 14, 2011]; Available from: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/infection-prevention-and-control-programs-in-ontario.html>.
11. Provincial Infectious Diseases Advisory Committee (PIDAC). Best Practices for Hand Hygiene in All Health Care Settings. 2010 [cited June 14, 2011]; Available from: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/hand-hygiene.html>.
12. Provincial Infectious Diseases Advisory Committee (PIDAC). Best Practices for Cleaning, Disinfection and Sterilization in All Health Care Settings. 2010 [cited June 14, 2011]; Available from: <http://www.oahpp.ca/resources/pidac-knowledge/best-practice-manuals/cleaning-disinfection-and-sterilization.html>.
13. Ontario. Ministry of Health and Long-Term Care. *Health Protection and Promotion Act: R.S.O. 1990*, chapter H.7. Toronto, Ontario 2008 [cited September 11, 2010]; Available from: http://www.e-laws.gov.on.ca/html/statutes/english/elaws_statutes_90h07_e.htm.

14. Ontario. Ministry of Health and Long-Term Care. Long-Term Care Homes Program Manual. [cited March 8, 2009]; 1-788]. Available from: http://www.health.gov.on.ca/english/providers/pub/manuals/ltc_homes/ltc_homes_mn.html#full.
15. Ontario. *Pesticides Act*, O. Reg. 63/09. 2009 [cited December 8, 2009]; Available from: <http://www.search.e-laws.gov.on.ca/en/isysquery/78220e09-8ed4-4861-9378-d81ecb148155/10/frame/?search=browseStatutes&context=>.
16. Centers for Disease Control and Prevention. Workbook for Designing, Implementing and Evaluating a Sharps Injury Prevention Program. Centers for Disease Control and Prevention.; 2004 [cited December 15, 2010]; 155]. Available from: <http://www.cdc.gov/sharpsafety/resources.html>.
17. Jernigan JA, Titus MG, Groschel DH, Getchell-White S, Farr BM. Effectiveness of contact isolation during a hospital outbreak of methicillin-resistant *Staphylococcus aureus*. *Am J Epidemiol*. 1996 Mar 1;143(5):496-504.
18. Siegel J, Rhinehart E, Jackson M, Chiarello L. The Healthcare Infection Control Practices Advisory Committee. Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings. *Am J Infect Control*. 2007 June, 2007;35(10 [Suppl 2]):S64-164.
19. Pittet D, Hugonnet S, Harbarth S, Mourouga P, Sauvan V, Touveneau S, et al. Effectiveness of a hospital-wide programme to improve compliance with hand hygiene. *Infection Control Programme*. *Lancet*. 2000 Oct 14;356(9238):1307-12.
20. Boyce JM. Strategies for controlling methicillin-resistant *Staphylococcus aureus* in hospitals. *J Chemother*. 1995 Jul;7 Suppl 3:81-5.
21. Conterno LO, Shymanski J, Ramotar K, Toyé B, Zvonar R, Roth V. Impact and cost of infection control measures to reduce nosocomial transmission of extended-spectrum beta-lactamase-producing organisms in a non-outbreak setting. *J Hosp Infect*. 2007 Apr;65(4):354-60.
22. Muto CA, Sistrom MG, Farr BM. Hand hygiene rates unaffected by installation of dispensers of a rapidly acting hand antiseptic. *Am J Infect Control*. 2000 Jun;28(3):273-6.
23. Whitby M, Pessoa-Silva CL, McLaws ML, Allegranzi B, Sax H, Larson E, et al. Behavioural considerations for hand hygiene practices: the basic building blocks. *J Hosp Infect*. 2007 Jan;65(1):1-8.
24. Harbarth S, Pittet D, Grady L, Goldmann DA. Compliance with hand hygiene practice in pediatric intensive care. *Pediatr Crit Care Med*. 2001 Oct;2(4):311-4.
25. Afif W, Huor P, Brassard P, Loo VG. Compliance with methicillin-resistant *Staphylococcus aureus* precautions in a teaching hospital. *Am J Infect Control*. 2002 Nov;30(7):430-3.
26. Moore D, Gamage B, Bryce E, Copes R, Yassi A. Protecting health care workers from SARS and other respiratory pathogens: organizational and individual factors that affect adherence to infection control guidelines. *Am J Infect Control*. 2005 Mar;33(2):88-96.
27. Kretzer EK, Larson EL. Behavioral interventions to improve infection control practices. *Am J Infect Control*. 1998 Jun;26(3):245-53.
28. Ontario. Ministry of Health and Long-Term Care. Infection Prevention and Control Core Competency Education Program. Routine Practices and Additional Precautions Module. 2007 [cited February 13, 2008]; Available from: http://www.health.gov.on.ca/english/providers/program/infectious/infect_prevent/ipc_cce_mn.html.
29. *Just Clean Your Hands*. Ontario's evidence-based hand hygiene program. Released 2008. [cited December 15, 2010]; Available from: <http://www.health.gov.on.ca/en/ms/handhygiene/>.
30. Vernon MO, Trick WE, Welbel SF, Peterson BJ, Weinstein RA. Adherence with hand hygiene: does number of sinks matter? *Infect Control Hosp Epidemiol*. 2003 Mar;24(3):224-5.
31. Berg DE, Hershov RC, Ramirez CA, Weinstein RA. Control of nosocomial infections in an intensive care unit in Guatemala City. *Clin Infect Dis*. 1995 Sep;21(3):588-93.

32. Pittet D. Improving compliance with hand hygiene in hospitals. *Infect Control Hosp Epidemiol.* 2000 Jun;21(6):381-6.
33. Picheansathian W. A systematic review on the effectiveness of alcohol-based solutions for hand hygiene. *Int J Nurs Pract.* 2004 Feb;10(1):3-9.
34. Boyce JM, Pittet D. Guideline for Hand Hygiene in Health-Care Settings. Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. *Infect Control Hosp Epidemiol.* 2002 Dec;23(12 Suppl):S3-40.
35. Kampf G, Kramer A. Epidemiologic background of hand hygiene and evaluation of the most important agents for scrubs and rubs. *Clin Microbiol Rev.* 2004 Oct;17(4):863-93.
36. Girou E, Loyeau S, Legrand P, Oppein F, Brun-Buisson C. Efficacy of handrubbing with alcohol based solution versus standard handwashing with antiseptic soap: randomised clinical trial. *BMJ.* 2002 Aug 17;325(7360):362.
37. Lewis AM, Gammon J, Hosein I. The pros and cons of isolation and containment. *J Hosp Infect.* 1999 Sep;43(1):19-23.
38. Kirkland KB, Weinstein JM. Adverse effects of contact isolation. *Lancet.* 1999 Oct 2;354(9185):1177-8.
39. Manian FA, Meyer L, Jenne J. Clostridium difficile contamination of blood pressure cuffs: a call for a closer look at gloving practices in the era of universal precautions. *Infect Control Hosp Epidemiol.* 1996 Mar;17(3):180-2.
40. Patterson JE, Vecchio J, Pantelick EL, Farrel P, Mazon D, Zervos MJ, et al. Association of contaminated gloves with transmission of Acinetobacter calcoaceticus var. anitratus in an intensive care unit. *Am J Med.* 1991 Nov;91(5):479-83.
41. Casanova L. Assessing the Risk of Viral Transmission from Contaminated Personal Protective Equipment to Employees' Skin and Clothing in the Healthcare Setting. 18th SHEA Annual Meeting; Orlando, Florida 2008.
42. Poutanen SM, Vearncombe M, McGeer AJ, Gardam M, Large G, Simor AE. Nosocomial acquisition of methicillin-resistant Staphylococcus aureus during an outbreak of severe acute respiratory syndrome. *Infect Control Hosp Epidemiol.* 2005 Feb;26(2):134-7.
43. Olsen RJ, Lynch P, Coyle MB, Cummings J, Bokete T, Stamm WE. Examination gloves as barriers to hand contamination in clinical practice. *JAMA.* 1993 Jul 21;270(3):350-3.
44. Doebbeling BN, Pfaller MA, Houston AK, Wenzel RP. Removal of nosocomial pathogens from the contaminated glove. Implications for glove reuse and handwashing. *Ann Intern Med.* 1988 Sep 1;109(5):394-8.
45. Health Canada. Infection Control Guidelines: Hand Washing, Cleaning, Disinfection and Sterilization in Health Care [currently under revision]. *Can Commun Dis Rep.* 1998 Dec;24 Suppl 8:1-55.
46. U.S. Food and Drug Administration. FDA Clears Glove Made from New Type of Latex. 2008 [cited October 22, 2008]; Available from: <http://www.fda.gov/bbs/topics/NEWS/2008/NEW01822.html>.
47. Wong TW, Lee CK, Tam W, Lau JT, Yu TS, Lui SF, et al. Cluster of SARS among medical students exposed to single patient, Hong Kong. *Emerg Infect Dis.* 2004 Feb;10(2):269-76.
48. Yu IT, Xie ZH, Tsoi KK, Chiu YL, Lok SW, Tang XP, et al. Why did outbreaks of severe acute respiratory syndrome occur in some hospital wards but not in others? *Clin Infect Dis.* 2007 Apr 15;44(8):1017-25.
49. Ulrich R, Quan, X., Zimring, C., Joesph, A, Quan X, Zimring C, Joseph A, Choudhary R. The Role of the Physical Environment in the Hospital of the 21st Century: A Once-in-a-Lifetime Opportunity. Report to The Center for Health Design for the *Designing the 21st Century Hospital* Project. 2004.
50. Bracco D, Dubois MJ, Bouali R, Eggimann P. Single rooms may help to prevent nosocomial bloodstream infection and cross-transmission of methicillin-resistant Staphylococcus aureus in intensive care units. *Intensive Care Med.* 2007 Mar 9.

51. Mulin B, Rouget C, Clement C, Bailly P, Julliot MC, Viel JF, et al. Association of private isolation rooms with ventilator-associated *Acinetobacter baumannii* pneumonia in a surgical intensive-care unit. *Infect Control Hosp Epidemiol*. 1997 Jul;18(7):499-503.
52. Chaudhury H. Advantages and Disadvantages of Single-Versus Multiple-Occupancy Rooms in Acute Care Environments. *Environment and Behavior*. 2005;37(6):760-86.
53. Rutala WA, Weber DJ. Surface disinfection: should we do it? *J Hosp Infect*. 2001 Aug;48 Suppl A:S64-8.
54. Sehulster L, Chinn RY. Guidelines for environmental infection control in health-care facilities. Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC). *MMWR Recomm Rep*. 2003 Jun 6;52(RR-10):1-42.
55. Smith PW, Rusnak PG. Infection prevention and control in the long-term-care facility. SHEA Long-Term-Care Committee and APIC Guidelines Committee. *Am J Infect Control*. 1997 Dec;25(6):488-512.
56. Grabsch EA, Burrell LJ, Padiglione A, O'Keeffe JM, Ballard S, Grayson ML. Risk of environmental and healthcare worker contamination with vancomycin-resistant enterococci during outpatient procedures and hemodialysis. *Infect Control Hosp Epidemiol*. 2006 Mar;27(3):287-93.
57. Bhalla A, Pultz NJ, Gries DM, Ray AJ, Eckstein EC, Aron DC, et al. Acquisition of nosocomial pathogens on hands after contact with environmental surfaces near hospitalized patients. *Infect Control Hosp Epidemiol*. 2004 Feb;25(2):164-7.
58. Huang SS, Datta R, Platt R. Risk of acquiring antibiotic-resistant bacteria from prior room occupants. *Arch Intern Med*. 2006 Oct 9;166(18):1945-51.
59. Hardy KJ, Oppenheim BA, Gossain S, Gao F, Hawkey PM. A study of the relationship between environmental contamination with methicillin-resistant *Staphylococcus aureus* (MRSA) and patients' acquisition of MRSA. *Infect Control Hosp Epidemiol*. 2006 Feb;27(2):127-32.
60. Rampling A, Wiseman S, Davis L, Hyett AP, Walbridge AN, Payne GC, et al. Evidence that hospital hygiene is important in the control of methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect*. 2001 Oct;49(2):109-16.
61. NHS Estates. The NHS Healthcare Cleaning Manual. 2007 [cited December 5]; Available from: http://patientexperience.nhsestates.gov.uk/clean_hospitals/ch_content/cleaning_manual/introduction.asp#manual.
62. Ontario. Ministry of Health and Long-Term Care. Ontario Regulation under the *Health Protection and Promotion Act* : Regulation 562 of R.R.O. 1990, Food premises, (as amended) Toronto, Ontario 2002 [cited December 8, 2009]; Available from: http://www.e-laws.gov.on.ca/Download?dDocName=elaws_regs_900562_e.
63. Ontario Hospital Association and the Ontario Medical Association Joint Communicable Diseases Surveillance Protocols Committee in collaboration with the Ministry of Health and Long-Term Care. Blood-Borne Diseases Surveillance Protocol for Ontario Hospitals. 2010 [cited December 15, 2010]; 1-22]. Available from: <http://www.oha.com/Services/HealthSafety/Documents/Protocols/Blood%20Borne%20Diseases%20Protocol%20-%20Reviewed%20and%20Revised%20November%202010.pdf>.
64. Ontario. *Occupational Health and Safety Act*. Ontario Regulation 474/07. Needle Safety. 2007 [cited December 8, 2009]; Available from: http://www.e-laws.gov.on.ca/Download?dDocName=elaws_regs_070474_e.
65. Scheckler WE, Brimhall D, Buck AS, Farr BM, Friedman C, Garibaldi RA, et al. Requirements for infrastructure and essential activities of infection control and epidemiology in hospitals: a consensus panel report. *Society for Healthcare Epidemiology of America. Infect Control Hosp Epidemiol*. 1998 Feb;19(2):114-24.
66. Gasink LB, Singer K, Fishman NO, Holmes WC, Weiner MG, Bilker WB, et al. Contact Isolation for Infection Control in Hospitalized Patients: Is Patient Satisfaction Affected? *Infect Control Hosp Epidemiol*. 2008 Mar;29(3):275-8.

67. Ontario Hospital Association and the Ontario Medical Association Joint Communicable Diseases Surveillance Protocols Committee in collaboration with the Ministry of Health and Long-Term Care. Introduction. 2010 [cited December 15, 2010]; 2]. Available from: <http://www.oha.com/Services/HealthSafety/Pages/CommunicableDiseaseBinder.aspx>.
68. Diekema DJ, Doebbeling BN. Employee health and infection control. *Infect Control Hosp Epidemiol*. 1995 May;16(5):292-301.
69. Nichol KL, Lind A, Margolis KL, Murdoch M, McFadden R, Hauge M, et al. The effectiveness of vaccination against influenza in healthy, working adults. *N Engl J Med*. 1995 Oct 5;333(14):889-93.
70. Ontario Hospital Association and the Ontario Medical Association Joint Communicable Diseases Surveillance Protocols Committee in collaboration with the Ministry of Health and Long-Term Care. Influenza Surveillance Protocol for Ontario Hospitals. 2010 [cited December 15, 2010]; 1-9]. Available from: <http://www.oha.com/Services/HealthSafety/Documents/Influenza%20Protocol%20Revised%20May%202010.pdf>.
71. Ontario Hospital Association and the Ontario Medical Association Joint Communicable Diseases Surveillance Protocols Committee in collaboration with the Ministry of Health and Long-Term Care. Rubella Surveillance Protocol for Ontario Hospitals. 2010 [cited December 15, 2010]; 1-9]. Available from: <http://www.oha.com/Services/HealthSafety/Documents/Rubella%20Protocol%20Revised%20May%202010.pdf>.
72. Ontario Hospital Association and the Ontario Medical Association Joint Communicable Diseases Surveillance Protocols Committee in collaboration with the Ministry of Health and Long-Term Care. Measles Surveillance Protocol for Ontario Hospitals. 2009 [cited December 15, 2010]; 1-9]. Available from: <http://www.oha.com/Services/HealthSafety/Documents/Protocols/Measles%20Protocol.pdf>.
73. National Advisory Committee on Immunization. Canadian immunization guide. 7th ed. [Ottawa]: Canadian Medical Association; 2006.
74. Ontario Hospital Association and the Ontario Medical Association Joint Communicable Diseases Surveillance Protocols Committee in collaboration with the Ministry of Health and Long-Term Care. Mumps Surveillance Protocol for Ontario Hospitals. 2009 [cited December 15, 2010]; 1-9]. Available from: <http://www.oha.com/Services/HealthSafety/Documents/Protocols/Mumps%20Protocol%20Revised%20January%202009.pdf>.
75. Ontario Hospital Association and the Ontario Medical Association Joint Communicable Diseases Surveillance Protocols Committee in collaboration with the Ministry of Health and Long-Term Care. Varicella/Zoster (Chickenpox/Shingles) Surveillance Protocol for Ontario Hospitals. 2010 [cited December 15, 2010]; 1-89]. Available from: <http://www.oha.com/Services/HealthSafety/Documents/Protocols/Varicella%20Protocol%20-%20Reviewed%20and%20Revised%20November%202010.pdf>.
76. Ontario Hospital Association and the Ontario Medical Association Joint Communicable Diseases Surveillance Protocols Committee in collaboration with the Ministry of Health and Long-Term Care. Pertussis Surveillance Protocol for Ontario Hospitals. 2009 [cited December 15, 2010]; 1-12]. Available from: <http://www.oha.com/Services/HealthSafety/Documents/Protocols/Pertussis%20Protocol%20Revised%20January%202009.pdf>.
77. Mulligan ME, Murray-Leisure KA, Ribner BS, Standiford HC, John JF, Korvick JA, et al. Methicillin-resistant *Staphylococcus aureus*: a consensus review of the microbiology, pathogenesis, and epidemiology with implications for prevention and management. *Am J Med*. 1993 Mar;94(3):313-28.
78. Boyce JM. Methicillin-resistant *Staphylococcus aureus* in hospitals and long-term care facilities: microbiology, epidemiology, and preventive measures. *Infect Control Hosp Epidemiol*. 1992 Dec;13(12):725-37.

79. Mayhall CG. Hospital Epidemiology and Infection Control. 3rd edition ed. Philadelphia: Lippincott Williams & Wilkins; 2004.
80. Sample ML, Gravel D, Oxley C, Toye B, Garber G, Ramotar K. An outbreak of vancomycin-resistant enterococci in a hematology-oncology unit: control by patient cohorting and terminal cleaning of the environment. *Infect Control Hosp Epidemiol*. 2002 Aug;23(8):468-70.
81. Jochimsen EM, Fish L, Manning K, Young S, Singer DA, Baker R, et al. Control of vancomycin-resistant enterococci at a community hospital: efficacy of patient and staff cohorting. *Infect Control Hosp Epidemiol*. 1999 Feb;20(2):106-9.
82. Austin DJ, Bonten MJ, Weinstein RA, Slaughter S, Anderson RM. Vancomycin-resistant enterococci in intensive-care hospital settings: transmission dynamics, persistence, and the impact of infection control programs. *Proc Natl Acad Sci U S A*. 1999 Jun 8;96(12):6908-13.
83. Macartney KK, Gorelick MH, Manning ML, Hodinka RL, Bell LM. Nosocomial respiratory syncytial virus infections: the cost-effectiveness and cost-benefit of infection control. *Pediatrics*. 2000 Sep;106(3):520-6.
84. Peel RK, Stolarek I, Elder AT. Is it time to stop searching for MRSA? Isolating patients with MRSA can have long term implications (letter). *BMJ*. 1997 Jul 5;315(7099):58.
85. Kellerman J, Rigler D, Siegel SE. The psychological effects of isolation in protected environments. *Am J Psychiatry*. 1977 May;134(5):563-5.
86. Stelfox HT, Bates DW, Redelmeier DA. Safety of patients isolated for infection control. *JAMA*. 2003 Oct 8;290(14):1899-905.
87. Johnson S, Gerding DN, Olson MM, Weiler MD, Hughes RA, Clabots CR, et al. Prospective, controlled study of vinyl glove use to interrupt *Clostridium difficile* nosocomial transmission. *Am J Med*. 1990 Feb;88(2):137-40.
88. Armstrong-Evans M, Litt M, McArthur MA, Willey B, Cann D, Liska S, et al. Control of transmission of vancomycin-resistant *Enterococcus faecium* in a long-term-care facility. *Infect Control Hosp Epidemiol*. 1999 May;20(5):312-7.
89. Tenorio AR, Badri SM, Sahgal NB, Hota B, Matushek M, Hayden MK, et al. Effectiveness of gloves in the prevention of hand carriage of vancomycin-resistant enterococcus species by health care workers after patient care. *Clin Infect Dis*. 2001 Mar 1;32(5):826-9.
90. Puzniak LA, Gillespie KN, Leet T, Kollef M, Mundy LM. A cost-benefit analysis of gown use in controlling vancomycin-resistant *Enterococcus* transmission: is it worth the price? *Infect Control Hosp Epidemiol*. 2004 May;25(5):418-24.
91. Slaughter S, Hayden MK, Nathan C, Hu TC, Rice T, Van Voorhis J, et al. A comparison of the effect of universal use of gloves and gowns with that of glove use alone on acquisition of vancomycin-resistant enterococci in a medical intensive care unit. *Ann Intern Med*. 1996 Sep 15;125(6):448-56.
92. Srinivasan A, Song X, Ross T, Merz W, Brower R, Perl TM. A prospective study to determine whether cover gowns in addition to gloves decrease nosocomial transmission of vancomycin-resistant enterococci in an intensive care unit. *Infect Control Hosp Epidemiol*. 2002 Aug;23(8):424-8.
93. Zachary KC, Bayne PS, Morrison VJ, Ford DS, Silver LC, Hooper DC. Contamination of gowns, gloves, and stethoscopes with vancomycin-resistant enterococci. *Infect Control Hosp Epidemiol*. 2001 Sep;22(9):560-4.
94. Xie X, Li Y, Chwang AT, Ho PL, Seto WH. How far droplets can move in indoor environments--revisiting the Wells evaporation-falling curve. *Indoor Air*. 2007 Jun;17(3):211-25.
95. Menzies D, Fanning A, Yuan L, Fitzgerald M. Tuberculosis among health care workers. *N Engl J Med*. 1995 Jan 12;332(2):92-8.
96. Public Health Agency of Canada. Canadian Tuberculosis Standards. 6th ed. Ottawa: Public Health Agency of Canada; 2007.

97. Canadian Standards Association. CAN/CSA Z317.2-01 (R2008) Special Requirements for Heating, Ventilation, and Air Conditioning (HVAC) Systems in Health Care Facilities. Toronto: Canadian Standards Association; 2001.
98. DeJoy DM, Murphy LR, Gershon RM. The influence of employee, job/task, and organizational factors on adherence to universal precautions among nurses. *Int J Ind Ergon.* 1995;16:43-55.
99. Clements A, Halton K, Graves N, Pettitt A, Morton A, Looke D, et al. Overcrowding and understaffing in modern health-care systems: key determinants in meticillin-resistant *Staphylococcus aureus* transmission. *Lancet Infect Dis.* 2008 Jul;8(7):427-34.
100. Health Canada. Guidelines for preventing the transmission of tuberculosis in Canadian Health Care Facilities and other institutional settings. *Can Commun Dis Rep.* 1996 Apr;22 Suppl 1:i-iv, 1-55.